



ANTI-FERTILITY ACTIVITY OF WHOLE PLANT EXTRACT OF *SARCOSTEMMA SECAMONE* (L) BENNET ON MALE ALBINO RATS

Thanga Krishna Kumari S¹, Sakthidevi G², Muthukumaraswamy S¹, Mohan VR^{2*}

¹Department of Botany, Sri K.G.S. Atrs College, Srivaikuntam, Tuticorin, Tamil Nadu, India

² Ethnopharmacology Unit, Research Department of Botany, V. O. Chidambaram College, Tuticorin, Tamil Nadu, India

Article Received on: 03/09/12 Revised on: 11/10/12 Approved for publication: 10/11/12

*Email vrmohanvoc@gmail.com

ABSTRACT

Anti-fertility effect of ethanol extract of whole plant extract of *Sarcostemma secamone* was observed in male albino rats. The relative weight of the testes and epididymis were decreased. The epididymal sperm count, motility and sperm abnormality were reduced significantly in treated rats. There was an increase in serum protein, albumin, globulin, urea, creatinine and the activity of liver marker enzymes (SGOT, SGPT and ALP) levels of control and treated rats. The activities of serum antioxidants (CAT, SOD, GPx, GST and GRD) in plant extract treated rats were decreased. The results of the hormonal assay showed that increased serum levels of FSH and estrogen but decreased in the serum levels of LH and testosterone compared to control. The results of fertility test indicated that the treated adult male rats reduced the number of female's impregnation. In addition, the number of implantations and the number of viable fetuses were also decreased. The results of the present study concluded that, ethanol extract of whole plant of *Sarcostemma secamone* inhibited sperm concentration, motility and testosterone which might result in a male fertility.

Key words: *Sarcostemma secamone*, Fertility, Testosterone, Antioxidant

INTRODUCTION

Population explosion is a leading cause of poverty and pollution in developing countries. Several potential approaches for infertility have been investigated over a long period, including chemical, hormonal and immunological approaches. However, no suitable method has emerged that is effective and free from side effects. The World Health Organization (WHO) has constituted a population control programme, which includes studies having traditional medicinal practices. Medicinal plants products have a long history of indigenous use in India as well as other countries¹. Phytotherapy has a very long tradition, although proper scientific explanation is relatively new. In our countries as well as in the world, there are several medicinal plants associated with anti-fertility properties²⁻⁴. Although very few contraceptives have been developed from plant extracts, their potentiality has not been determined accurately and their mode of action has been beyond our knowledge until now because there are many problems in assessing plant extract including batch to batch variation and a lack of definite active portion of the extract used for the development of herbal contraceptives. Fertility regulation with plants or plant preparations has been reported in the ancient literature of Indigenous systems of medicine. A large number of plant species with anti-fertility effects have been screened in China and India beginning about 50 years ago and were subsequently fortified by National and International agencies^{5,6}. However, the search for an orally active, safe and effective plant preparation or its compound is yet to be needed for fertility regulation due to incomplete inhibition of fertility or side effects.

Sarcostemma secamone (L) Bennet is an important medicinal plant belonging to the family Asclepiadaceae. It is used in the traditional systems of medicine for various ailments. The decoction of the plant is useful to gargle for throat and mouth infection. The latex is bitter and used as a vulnerary. Fresh roots are prescribed for jaundice^{7,8,9,10}. The milky sap forms a wash for ulcers. In combination with turpentine, it is prescribed for itch¹¹. The plant is hot, bitter, tonic, expectorant, pungent, dry and indigestible causes flatulence,

diuretic, laxative, aphrodisiac, anthelmintic, useful in leucoderma and bronchitis. The juice is used in gleet, gonorrhoea, pain in the muscles, cough and given to children as an astringent¹². Leaf powder stimulates arculatory system, increases secretion of urine and activates uterus¹³. The fruit juice is used in gonorrhoea and pain in muscles. The leaves, roots and latex of *Sarcostemma secamone* are employed in treating many diseases like mouth ulcer, sour throat, jaundice and ulcers¹⁴⁻¹⁶. In view of the above said medicinal properties, the present study was carried out to test the efficacy of anti-fertility activity of ethanol extract of whole plant of *Sarcostemma secamone* on male albino rats.

MATERIALS AND METHODS

Plant material

The well grown whole plant of *Sarcostemma secamone* (L.) Bennet was collected from natural forests of Western Ghats at Thanniparai, Srivilliputhur, Virudhunagar District, Tamil Nadu. The collected plants were identified by the Botanical Survey of India, Coimbatore. A voucher specimen was retained in Ethnopharmacology unit, Research Department of Botany, V.O.Chidambaram College, Tuticorin for further reference.

Preparation of plant extract

The whole plant of *Sarcostemma secamone* were dried separately under shade and then powdered with a mechanical grinder to obtain a coarse powder, which were then subjected to extraction in a Soxhlet apparatus using ethanol. The ethanol extract were concentrated in a rotatory evaporator. The concentrated ethanol extract of whole plant of *Sarcostemma secamone* were used for anti-fertility activity.

Animals

Normal healthy male Wistar albino rats (180-240g) were used for the present investigation. Animals were housed under standard environmental conditions at temperature (25±2⁰C) and light and dark (12:12h). Rats were fed with standard pellet diet (Goldmohur brand, MS Hindustan Lever Ltd., Mumbai, India) and water *ad libitum*.

Acute Toxicity Studies

Acute oral toxicity study was performed as per OECD-423 guidelines (acute toxic class method), albino rats of either sex selected by random sampling were used for acute toxicity study¹⁷. The animals were kept fasting for overnight and provided only with water, after which the extracts were administered orally at 5mg/kg body weight by gastric incubations and observed for 14 days. If mortality was observed in two out of three animals, then the dose administered was assigned as toxic dose. If mortality was observed in one animal then the same dose was repeated again to confirm the toxic dose. If mortality was not observed, the procedure was repeated for higher doses such as 50, 100 and 2000 mg/kg body weight.

Experimental Design

The male rats were divided into 3 groups consisting of 5 animals.

Group I: Rats received normal saline daily for 14 days, orally. (Normal control).

Group II: Rats received ethanol extract of whole plant of *Sarcostemma secamone* at the dose of 150 mg/kg body weight daily for 14 days.

Group III: Rats received ethanol extract of whole plant of *Sarcostemma secamone*, at the dose of 300 mg/kg body weight daily for 14 days.

After 24 hours of last treatment, the final weight was recorded and the animals were sacrificed by decapitation. Blood was collected. Sera were separated by centrifugation at 3000g for 10 minutes and stored at 20°C until used for various biochemical assays. Then testes, epididymis, vas deferens, seminal vesicle and ventral prostate were dissected out, trimmed off extraneous and weighed accurately on torsion balance. The organs weights were expressed in terms of mg/100g body weight.

Sperm count

Epididymal fluid (for sperm count) was collected from caput and cauda segments separately and diluted with Sorenson's buffer (pH 7.2). The separated fluid was taken for sperm count. Sperm count was carried out by using Neubauer's haemocytometer as described by Zaneveld and Pelakoski¹⁸.

Sperm motility and abnormality

After anaesthetizing the rats, the caudal epididymis was then dissected. An incision (about 1mm) was made in the caudal epididymis and drops of sperm fluid were squeezed onto the

microscope slide and 2 drops of normal saline were added to mobilize the sperm cells. Epididymal sperm motility was then assessed by calculating motile spermatozoa per unit area.

Morphology (abnormality) was evaluated on sperm from the caudal epididymis. The total morphological abnormalities were observed as described by Linde *et al*¹⁹.

Serum biochemical analysis

Serum proteins²⁰ and serum albumins were determined by quantitative colorimetric method by using bromocresol green. The total protein minus albumin gives the globulin, urea²¹, creatinine²², serum glutamate pyruvate transaminase (SGPT) and serum glutamate oxaloacetate transaminase (SGOT) was measured spectrophotometrically by using the method of Reitman and Frankel²³. Serum alkaline phosphatase (ALP) was measured by the method of King and Armstrong²⁴.

Serum antioxidants

Serum antioxidant Catalase (CAT)²⁵, Superoxidedismutase (SOD)²⁶, Glutathione peroxidase (GPx)²⁷, Glutathione s-transferase (GST)²⁸ and Glutathione reductase (GRD)²⁹ were analyzed.

Hormonal Assay

Blood removed from the rats by intracardiac method. Blood was centrifuged at 3000 rpm to separate the serum for the measurement of testosterone, Luteinizing hormone (LH), estrogen and follicle stimulating hormone (FSH). The quantitative determination of hormones was done by using Enzyme Immunoassay Method (EIA). The EIA kit was obtained from Immunometrics (London, UK).

Fertility test

Fertility was estimated in adult male rats treated with ethanol extracts of whole plant of *Sarcostemma secamone* and in the control males counterparts. Each male was placed in an individual cage with two virgin untreated females of the same strain. They were left together for 10 days during which two estron cycles had elapsed. One week after the removal of the exposed males, pregnant females were killed by cervical dislocation under light ether anesthesia and the number of implantation sites, the number of fetuses and the number of resorption sites were recorded³⁰.

Statistical Analysis

Data were expressed as Mean ± SEM. Student's t test was used for statistical comparison.

Table 1: Effect of *Sarcostemma secamone* whole plant extracts on the body and reproductive organ weight of adult male albino rats

Treatment Groups	Body wt (gm)		Testis (gm)	Epididymis (mg)		VD (mg)	SV (mg)	Prostrate (mg)
	Before	After		Caput	Cauda			
Group-I	211±12.4	204±11.6	1.929±1.1	143.61±2.5	256.56±12.4	294.51±12.98	405.56±15.46	178.31±11.33
Group-II	198.48±8.56	161.33±9.56*	1.629±0.84*	139.59±8.11	162.66±7.98*	251.33±8.24*	293.13±10.47*	173.29±11.24
Group-III	253.17±13.38	210.69±11.39	1.688±0.92*	130.11±6.83	144.39±8.53*	249.50±9.56*	286.26±11.36*	181.56±10.14

Each value is SEM of 5 animals *P<0.05, Control Vs Treated

Table 2: Effect of *Sarcostemma secamone* whole plant extracts on the sperm concentration and motility in the epididymis of adult male albino rats

Treatment Groups	Sperm concentration (counts × 10 ⁶ mil)		Sperm motility (FMI)@(cauda)	Sperm abnormality #	
	caput	cauda		Head (%)	Tail (%)
Group-I	445.63±6.84	521.39±8.36	114.21±2.56	8.31±1.2	5.66±0.94
Group-II	326.54±9.33**	349.51±10.27**	58.14±1.26**	41.55±2.53*	53.26±2.04**
Group-III	304.67±8.26**	321.68±8.39**	62.11±1.07**	52.16±2.27**	61.59±1.14**

Each value is SEM of 5 animals *P<0.05; **P<0.01, Control Vs Treated.

@ : Motility is movement recorded after 5 min in suspension of caudal epididymal spermatozoa in phosphate buffered solution

: Expressed in percentage

Table 3: Effect of *Sarcostemma secamone* whole plant extracts on few serum biochemical profile of adult male albino rats

Parameter	Group-I	Group-II	Group-III
Protein (gm/dl)	8.22±0.24	7.13±0.33*	7.29±0.53*
Albumin (gm/dl)	1.54±0.11	4.03±0.34	3.93±0.19
Globulin (gm/dl)	3.68±0.34	3.10±0.34	3.36±0.18
A/G Ratio:	1:2:1	1:3:1	1:2:1
Urea (mg/ml)	13.27±1.26	16.27±1.98	26.56±2.31
Creatinine (mg/dl)	0.71±0.03	1.26±0.10	1.59±0.19
SGOT (U/L)	11.37±1.13	15.56±1.04	14.31±1.34
SGPT (U/L)	24.63±1.28	23.14±1.16	27.33±1.11
ALP (U/L)	163.27±5.6	163.31±4.56	173.39±5.86

Each value is SEM of 5 animals * $P<0.05$, Control Vs Treated

Table 4: Effect of *Sarcostemma secamone* whole plant extracts on the activity of serum Catalase, Glutathione peroxidase, Glutathione-S-transferase, Superoxide dismutase and Glutathione reductase in rats

Treatment	Catalase (μ moles of H_2O_2 decomposed/min/mg protein)	Glutathione peroxidase (μ moles of NADPH oxidized/min/mg protein)	Glutathione-S-transferase (μ moles of conjugate formed/min/mg protein)	Superoxide dismutase (Units)	Glutathione reductase (μ moles of NADPH oxidized/min/mg protein)
Group-I	7.56±0.51	0.304±0.05	11.59±1.08	20.36±1.14	29.33±1.65
Group-II	3.24±0.12**	0.211±0.05**	6.91±0.56*	11.24±1.24**	15.33±1.22*
Group-III	3.01±0.16**	0.183±0.02***	3.54±0.14**	7.31±1.56***	11.46±1.65*

Values are given as means \pm S.D from six rats in each group * $P<0.05$; ** $P<0.01$; *** $P<0.001$ Control Vs Treated

Table 5: Effect of *Sarcostemma secamone* whole plant extracts on sex hormones levels and pituitary gonadotrophins in male albino rats

Treatment Groups	Parameters			
	Testosterone (mg/ml)	LH/ICSH (μ lu/ml)	Estrogen (pg/ml)	FSH (μ lu/ml)
Group-I	4.39±0.17	3.93±0.26	21.33±1.05	1.17±0.31
Group-II	2.16±0.16**	1.73±0.16**	38.33±1.39**	2.86±0.15
Group-III	1.84±0.24**	1.21±0.11**	40.14±1.26**	2.99±0.21*

Each value is SEM 5 animals * $P<0.05$; ** $P<0.01$, Control Vs Treated

Table 6: Effect of *Sarcostemma secamone* whole plant extracts on the fertility of male albino rats

Groups	No. of male	No. of females	No. of pregnant females	No. of implantation	No. of viable fetuses	Total No. of resorption sites
Group-I	4	8	8/8(100%)	10.21±1.06	4.35±1.14	3
Group-II	4	8	2/8(25%)**	6.23±1.54*	3.06±1.12*	6
Group-III	4	10	3/10(30%)**	4.01±1.31**	2.84±1.23**	5

Each value is SEM 5 animals * $P<0.05$; ** $P<0.01$, Control Vs Treated

RESULTS

Body and Reproductive organ weight

The ethanol extract of whole plant of *Sarcostemma secamone* at different concentration were treated on male Wistar albino rats for anti-fertility activity.

The administration of ethanol extract of whole plant of *Sarcostemma secamone* to rats, slightly decrease the body weight (Table-1) and on the libido of treated rats whereas weight of testes and other accessory organs were decreased ($P<0.01$) (Table-1). Among the accessory sex organs, a significant weight reduction was seen in the caput and caudal epididymal segment. The weight reduction was dose-dependent i.e. high dose (300mg/kg body weight) treated groups (Group-III) drastically reduced followed by less in

low dose group (150mg/kg body weight) (Group-II). Slight changes were observed in vasdeferenes, seminal vesicle and prostate.

Sperm count and Sperm motility

Sperm motility and sperm density in caudal epididymis, significantly decreased and the reduction was severe in higher dose treated group (Group-III) followed by low dose group (Group-II) (Table-2) and the same trend was seen in the caput epididymal sperm density when compared to control (Group-I).

Serum biochemical profile

Serum protein, albumin, globulin, urea, creatinine and the activity of liver marker enzymes (SGOT, SGPT and ALP) levels of control and treated rats were depicted in the Table-3.

Except protein albumin and globulin, all the other parameters were slightly increased.

Serum antioxidants

The activities of CAT, SOD, GPx, GST and GRD in the serum of control and plant extract treated rats were presented in Table-4. In the present study, plant extract treated rats had shown decreased activities of all the studied antioxidants when compared to control rat.

Reproductive hormone level

Serum Testosterone level

The ethanol extract of whole plant of *Sarcostemma secamone* (150 and 300 mg/kg body weight) repeated treatment for 14 days caused a significant decrease in serum level of testosterone in male rats. The level of testosterone decrease was dose related (Table-5).

Serum luteinizing hormone (LH) level

Repeated treatment of male rats with the ethanol extract of whole plant of *Sarcostemma secamone* for 14 days caused a dose related decrease in the serum level of LH. The level of decrease is statistically significant.

Serum Estrogen level

The ethanol extract of whole plant of *Sarcostemma secamone* (150 and 300 mg/kg body weight) caused an increase in the serum level of estrogen in male rats. Doses of 150 and 300 mg/kg body weight administered daily for 14 days cause raise in the serum level of estrogen.

Serum Follicle Stimulating Hormone (FSH) level

Pre-treatment with ethanol extract of whole plant of *Sarcostemma secamone* caused an increase in the serum level of FSH male rats compared to control. An increase in the serum level of FSH was statistically significant.

Fertility test

The results represented in Table-6 shows that intragastric administration of ethanol extract of whole plant of *Sarcostemma secamone* (150 and 300 mg/kg body weight) for 14 days to male rats causes a significant decrease ($P < 0.05$) in the number of females impregnated by male treated rats. The number of viable fetuses calculated after cesarean sections were significantly decreased ($P < 0.05$) in female rats impregnated with untreated male rats. On the other hand, the number of resorption sites was found to be increased in female impregnated by extract treated male rats when compared to controls.

DISCUSSION

Studies on the effects of plant products on male reproductive system and fertility are comparatively few and far fetched. From a public health perspective, the head for contraception has never been greater.

The results revealed slight changes in the body weight of rats treated with ethanol extract of whole plant of *Sarcostemma secamone* (150 and 300 mg/kg body weight) for 14 days. The testes and other accessory sex organs, a significant weight reduction were seen in the testes, caput and caudal epididymal segments and the weight reduction was dose dependent. Reduction in the weight of testes and other accessory sex organs might be due to low level of androgen, which was not enough to maintain the weight of gonads and accessories³¹. It is known that the accessory sex organs viz., epididymis and vasdeferens are androgen dependent target organs and manifest differential sensibility to androgens for maintenance of their structure and function. It is also known that, any change in circulating androgens would affect the internal microenvironment of epididymis and thereby lead to alteration in sperm motility and metabolism³².

In the present study, ethanol extract of whole plant of *Sarcostemma secamone* treated rats decreased the sperm motility and sperm density in caudal and caput epididymal segments (Table-2). Drastic effect on the nature of the normal sperms in the caput and caudal region was observed in ethanol extract of whole plant of *Sarcostemma secamone* treated rats. Further tail region of the sperm in all the treated groups (Group-II & Group-III) were much affected than the head regions (Table-2). The development of normal and mature sperm is the key to optimum male fertility. The production of the sperm cells (spermatozoa) and testosterone in the testes are mainly regulated by the follicle stimulating hormone (FSH) and luteinizing hormone (LH), which are released from the anterior pituitary³³. FSH stimulates spermatogenesis in the steroli cells, while LH simulates the production of testosterone in the leydig cells of the testes³⁴. Many studies on the testes of rat treated with plant extracts has also demonstrated that the inhibitory activity on the proliferation of spermatogonia in mammals³⁵⁻³⁷. Spermatogenesis is therefore, a complicated process, covering proliferation of the spermatogonia, long-lasting process of the tissue meiosis and numerous changes in the spermatids during their pre-formation. The result of the present study suggest that ethanol extract of whole plant of *Sarcostemma secamone* for 14 days may affect the normal function of the steroli and leydig cells.

Sexual cells can occur during the reproductive phase, mitotic division of the spermatogonia or during the maturation of the spermatozoa, thereby affecting the number and quality of the sperm cells produced in the testes. Among the ethanol extract treated groups II and III (150 and 300mg/kg body weight) produced a significant reduction in the sperm count and viable sperms. This may be as a result of the ability of the extract at the given doses, to either interfere with spermatogenetic process in the somniferous tubules, epididymal functions or activities of testosterone on hypothalamic release factor and anterior pituitary secretion of gonadotropins which may result in alteration of spermatogenesis^{38, 39}. The presence of immature sperms was also observed in the experimental rats treated with ethanol extract of whole plant of *Sarcostemma secamone* (150 and 300 mg/kg body weight). This suggests that the 150 mg/kg body weight and 300 mg/kg body weight dose level could affect the maturation of the spermatozoa in the male rats, which might also be a contributory factor to the decrease in the mean total sperm count. The data generated in the present study, by and large, confirm to those already reported and studied with various plant extracts⁴⁰⁻⁴². The decrease in the caudal epididymal sperm counts are clear indications that, the ethanol extract of whole plant of *Sarcostemma secamone* can affect one or more aspects of spermatogenesis as well as spermigenesis. Though a direct effect of the ethanol extract of whole plant of *Sarcostemma secamone* on the cellular mechanisms of spermatogenesis cannot be concluded, it is likely that the impairment of the hormonal mechanisms concerned with the regulation of spermatogenesis may be the underlying cause.

The various other sperm abnormalities like sluggish motility, coiled tail and sperm maturation are also due to *Sarcostemma secamone* toxicity. The hitherto unreported abnormal sperm methodology, coiled tail and malformed head could be attributed to both testicular and epididymal effects of the ethanol extract of whole plant of *Sarcostemma secamone*. Coiling of the sperm tail is usually the product of abnormal axoneme and/or the other dense fibril. The outcome of the

present study affirms the male reproductive toxic effects of *Sarcostemma secamone* when applied as a therapeutic agent. Since male reproductive toxicology and male contraception are two sides of the same coin, the negative consequence of *Sarcostemma secamone* on the sperm may be taken as an advantage for further study. By the treatment employed in the study, no toxic effect was produced in the liver and kidney, neither was it directly involved on the development and functioning of the male reproductive system nor in the reproductive organs.

Superoxide dismutase (SOD) scavenges both extracellular and intracellular superoxide anion and prevents lipid peroxidation of the plasma membrane. In order to act against H_2O_2 it must be conjugated with catalase or glutathione peroxidase⁴³. The reduced level of catalase, glutathione peroxidase, glutathione s-transferase and glutathione reductase might be due to the excess production of anions in response to the ethanol extract of whole plant extract of *Sarcostemma secamone*. It is possible that an increased rate of ROS production may inhibit the action of these antioxidant enzymes or alternatively the decreased expression of these antioxidant enzymes may cause increased oxidative stress⁴⁴. This will result in increased LPO, decreased sperm motility, viability and function, an ultimately leads to infertility.

The present study revealed a decrease in the serum level of testosterone. This observation was similar to the earlier findings of⁴⁵⁻⁴⁷. The reduction in the serum level of testosterone could probably be due to the decrease of serum levels of LH/ICSH observed in this investigation. Leydig cells secrete testosterone by the stimulatory effect of LH^{48, 49}. In males reduction of testosterone level may impair spermatogenesis and cause male infertility. This study further observed a dose dependent increase in the serum estrogen level. This increase might probably be due to the conversion of testosterone to estrogen^{50, 51}.

Treatment with the ethanol extract of whole plant of *Sarcostemma secamone* (150 and 300mg/kg body weight) was highly effective in producing reversible functional sterility. The drug treated male rats clearly indicated structural and functional alteration in testes, epididymis and seminal vesicle. Depletion of sperm count and sperm motility in the drug treated rats suggests alteration in sperm motility resulted in partial infertility within 14 days. This resulted in abnormal sperm functions which ultimately gave rise to complete male sterility. Among the plant based contraceptives, inhibition of male fertility after administration of natural substances has been related to decreased spermatozoa density⁵². For male contraception, it is not necessary to stop spermatogenesis, but it is enough to eliminate the fertilizing ability of the spermatozoa by causing changes in the morphology or in the function of the sperm⁵³. In the present study, a significant decrease in the sperm density and motility was observed in the cauda epididymis in the entire treatment group, which lead to proven in the impairment of fertility in all the treated groups. The results presented in this study also indicated that the treatment with the ethanol extract of whole plant of *Sarcostemma secamone* by adult male rats reduces the number of female's impregnation. In addition, the number of implantations and the number of viable fetuses were also decreased, this could be a reflect and may be due to the decrease in sperm motility and sperm density observed in this study. Hence, this may be due to the effects of the given plant extracts on the enzymes involved in the oxidative phosphorylation.

From the present study it can be concluded that *Sarcostemma secamone* is capable to suppress male fertility without altering general metabolism. Hence the possible male contraceptive efficacy of *Sarcostemma secamone* whole plant extract cannot be ignored paving way to the smooth development for the clinicians' interests in clinical trials towards emergence of a potent herbal male contraceptive.

Recently many laboratories are engaged in developing male contraceptives from plants⁵⁴. Plant products as contraceptives will be more acceptable for economic reasons in terms of self reliance and the possible practicability for a male pill approach in countries where population pressure is high. Recently extensive effects have been made to study the anti-fertility drugs from plants^{55,56}. In the present study, dose dependent treatment of *Sarcostemma secamone* whole plant extract and duration suggests marked alterations in the male reproductive organs. Further studies are needed to prove whether the alterations are reversible or permanent after cessation of treatment and for understanding the extract mechanism.

ACKNOWLEDGEMENT

The authors are thankful to Dr.R.Sampathraj, Honorary Director, Dr.Samsun Clinical Research Laboratory, Tirupur for providing necessary facilities to carry out this work

REFERENCES

- Hinemath SP, Badani S, Hunasagatta SK and Patil SB. Anti-fertility and hormonal property of flavones of *Striga oroleanchoider*. Eur J Pharmacol 2000; 391:193-7.
- Madhumita G and Saral AM. Free radical scavenging assay of theyetia nerrifolia leaf extracts. Asaian Journal of Chemistry 2009; 21:2468-2470.
- Madhumita G and Saral AM. Antimicrobial activity of successive extracts of *Thevetia nerrifolia*. Asaian Journal of Chemistry 2009; 21:2471-2472.
- Hinemath SP, Rudresh K, Badani S, Hunasagatta SK, Patil SB and Patil SR. Post-coital anti-fertility of *Acalypha indica*. L. J Ethnopharmacol 1999; 67:226-8.
- WHO. Reproductive health research at WHO: a new beginning Biennial Report 1998-99, Special Programme of Research, Development and Research Training in Human Reproduction, World Health Organization, Geneva, 2000.
- Lohiya NK. Plant products for contraception: how to make it a reality? In: Puri, CP (Edn.), ISSRF News letter, Indian society for the study of reproduction and fertility, Mumbai. 2000; 5:9-12.
- Chopra RN, Nayar SL and Chopra IC. Glossary of Indian Medicinal plants. CSIR, New Delhi, India. 1956.
- Chopra RN, Chopra IC, Handa KL and Kapoor LD. Indigenous Drugs of India. Academic publishers, Calcutta. India. 1958.
- Anonymous. The wealth of India- A dictionary of Indian raw materials and industrial products. Vol-Vii (N-Pe) CSIR, New Delhi, India. 1966.
- Nadkarni AK. Indian material medica, popular prakashan pvt Ltd., Mumbai. 1982.
- Kritikar KR and Bassu BD. Indian medicinal plants, International book Distributors, Dehradun, India. 1976.
- Poornima N, Umarraian KM and Babu K. Studies on Anatomical and phytochemical Analysis of *Oxystelma esculentum* (L.f) R. Br. Ex Schlt. J Bot Res Internatoinal. 2009; 2:239-243.
- Prajapati ND, Purohit SS, Sharma AK and Kumar T. A Hand book of medicinal plants, Agrobios, India.
- Khan AV. Thesis submitted to Aligarh muslim university, Aligarh. 2002.
- Sathyavathi GV, Gupta AK and Janabu N. Medicinal plants of India, CSIR publication, Indian council of Medical Research, Cambridge, New Delhi. 1987; 2: X1 + 557.
- Jain SK. Dictionary of Indian folk medicine and ethnobotany, Deep publication, New Delhi. 1991; pp.X11+311.
- OECD. (Organization for Economic Cooperation and Development). OECD guidelines for the testing of chemicals/section 4: Health Effects Test No.423; Acute Oral Toxicity-Acute Toxic class method, OECD.
- Zaneveld LJD and Pelakoski. Collection and physical examination of the ejaculate. In: Hafez ESE(ed). Techniques in human andrology. Vol.I,

- Human reproductive medicine. North-Holland Publishing company, Asterdam, 1997, pp. 147-172
19. Linde RE, Strader LF, Slot VL and Suarez JD. End points of spermatotoxicity in the rat after short duration exposures to fourteen reproductive toxicants. *Reprod. Toxicol.* 1992; 6: 491-505.
 20. Lowry OH, Rosenbrough NJ, Farr AL, Randall RJ. Protein measurement with the folin's phenol reagent. *Journal of Biological Chemistry.* 1951; 193: 265-275.
 21. Varley H. *Practical clinical biochemistry*, Arnold Heinemann Publication Pvt. Ltd. 1976, pp. 452.
 22. Owen JA, Iggo JB, Scongrett FJ and steward IP. Determination of creatinine in plasma serum, a critical examination. *J.Biochem.* 1954; 58: 426-437
 23. Reitman S and Frankel SA. Colorimetric method for the determination of serum glutamic oxaloacetic and glutamic pyruvic transaminase. *Am.J.Clin. Pathol.* 1957; 28: 56-63.
 24. King EJ and Armstrong AR. Determination of serum and bile phosphate activity. *Can. Med. Assoc. J.* 1934; 31: 56-63.
 25. Sinha AK. Colorimetric assay of catalase. *Anal Biochem* 1972; 47: 389-394.
 26. Das S, Vasight S, Snehlata R, Das N and Srivastava LM. Correlation between total antioxidant status and lipid peroxidation in hypercholesterolemia. *Curr Sci.* 2000; 78: 486-487.
 27. Rotruck JT, Pope AL, Ganther HE and Swanson AB. Selenium: Biochemical roles as a component of Glutathione peroxidase. *Science* 1984; 179: 588-590.
 28. Habig WH, Pabst MJ and Jaco WB. Glutathione s-transferase. The first enzymatic step in mercapturic acid formation. *J Biol Chem* 1974; 249: 7130-7139.
 29. Goldberg DM and Spooner RJ. Glutathione reductase In: *Methods in enzymatic analysis*, V.C.H. Weinheim, Germany. 1983; 258-265.
 30. Rugh R. *The mouse, its reproduction and development.* Burgess, Minneacolis. 1968.
 31. Sharma N and Jacob D. Anti-fertility investigation and toxicological screening of the Petroleum ether extract of the leaves of *Mentha arvensis* L. in male albino mice. *J Ethanopharmacol* 2001; 75: 5-12.
 32. Khan PK and Awasthy KS. Cytogenetic toxicity of Neem. *Food chem. Toxicol* 2003; 41: 1325-1328.
 33. Steinberger E. Hormonal control of mammalian Spermatogenesis. *Physiology Review* 1971; 5111, 1-22.
 34. Kerr JB and Klester DM. Cyclic variation in steroli cell lipid content throughout the spermatogenic cycle in the rats. *J. Reprod Fert* 1975; 43: 1-8.
 35. Steinberger E, Steinberger A and Perlof WH. Initiation of spermatogenesis *In vitro*. *Endocrinology* 1964; 74: 788.
 36. Mancini RE, Castro A and Serguer AC. Histological localization of follicle-stimulating and luteinizing hormones in the rat testes. *J Histochem. Cytochem* 1967; 15: 516-526.
 37. Krueger PM, Hodgen CD and Sherins KI. New evidence for the role of the steroli cells and spermatogonia in feed back control of FSH secretion in male rat. *Endocrinology* 1974; 95: 955-962.
 38. Bowman WCM and Rand MJ. The reproductive system and drugs affecting the reproductive systems. *Text book of Pharmacology* 1985; 2.
 39. William KN. Hormones and Hormone antagonists. In: *Remington, The Science and Practice of Pharmacy* 2000; 77: 1390-1391.
 40. Njar vc, Alao TO, Okogun JI, Raji Y, Bolarinwa AF and Nduka EV. Anti-fertility activity of *Quassia amara*: quassin inhibits the steroidogenesis in rat Leydig cells *in vitro*. *Planta Med* 1995; 61: 180-182.
 41. Raji Y and Bolarinwa AF. Anti-fertility activity of *Quassia amara* in male rats- *in vivo* study. *Life Sci* 1997; 61: 1067-1074.
 42. Parveen S, Das S, Kundra CP and Pereira BM. Acomprehensive evaluation of the reproductive toxicity of *Quassia amara* in male rats. *Reprod. Toxicol* 2002; 17: 45-50.
 43. Jeulin C, Soufir JC, Weber P, Laval-Martin D and Calvayrac R. Catalase activity in human spermatozoa and seminal plasma. *Gamete Res* 1989; 24: 185-196.
 44. Sikka SC, Rajasekaran M and Hellstrom WJ. Role of oxidative stress and antioxidants in male infertility. *J. Androl* 1995; 16: 464-8.
 45. Udoh P and Kehinde A. Studies on anti-fertility effects of paw paw seeds (*Carica papaya*) on the gonads of male Albino rats. *Phytother. Res* 1999; 13: 226-228.
 46. Udoh P and Ekipeyong J. Effects of *Mucuna urens* (Horse eye bean) on the gonads of male Guinea pigs. *Phytother. Res* 2001; 15: 99-102.
 47. Udoh P, Essien I and Udoh F. Effect of *Carica papaya* (paw paw) seeds extract on the morphology of pituitary-gonadal axis of male albino rats. *Phytotherapy. Res* 2005 a; 19: 1065-1068.
 48. Udoh FV and Udoh PB. Hepatotoxicity of methanol extract of *Carica papaya* seeds in Wistar rats. *Pharmaceutical Biol* 2005; 43: 349-352.
 49. Udoh FV, Udoh PB and Umon EE. Activity of alkaloid extract of *Carica papaya* seeds on reproductive functions in male Wistar rats. *Pharmaceutical Biol* 2005 b; 43: 563-567.
 50. Carr BR and Blackwell RE. Neuroendocrinology of reproduction In: *Text book Reproductive Medicine* Carr and Blackwell RE (eds). Norwalk: Appleton and Lange. 1993; 157.
 51. Chinoy RJ and Padman P. anti-fertility investigation and benzene extract or *Carica papaya* seeds in male albino rats. *J Med Aromatic Plant Sci* 1996; 18: 489-494.
 52. Witche P, Kamtechouing P, Sokeng S, Moundipa PF, Tantchu J, Essame JL and Kaveeta N. Reversible antispermatogenic and anti-fertility activity of *Mondia whitei* L. in male albino rats. *Phytother. Res* 2001; 15: 26-9.
 53. Dwivedi AK, Chaudhary M and Sarine J.Ps. standardization of a new spermicidal agent sapindus saponin and its estimation in its formulation. *Indian J. Pharm. Sci* 1990; 52: 165-167.
 54. US National Academy of Sciences, Publication. *Neem-a tree for solving global problems.* For thousands of years the beneficial properties of Neem office of International Affairs. 1992.
 55. Handelsman DJ. Hormonal male contraception. In: Puri CP, Van Lock PFA. Editors. *Current concepts in fertility regulation and reproduction.* New Delhi. Wiley Eastern Ltd. 1994; 133.
 56. Upadhyay SN, Dhawan S and Talwar CP. Anti-fertility effects of Neem (*Azadirachta indica*) oil in male rats by single intra-vas administration. An alternate approach to vasectomy. *J. Androl* 1993; 14: 275-281.

Source of support: Nil, Conflict of interest: None Declared