



Research Article

ANALYSIS OF VOLATILE COMPOUNDS IN THE SAP OF *AZADIRACHTA INDICA* (NEEM) USING GAS CHROMATOGRAPHY MASS SPECTROMETRY

Praveen Kumar P¹, Nandhini A R¹, Subhapiya P² and Gowri Shankar B A^{1*}

¹Department of Biotechnology, Laboratory for Venom Peptidomics and Molecular Simulation, Bannari Amman Institute of Technology, Erode, Tamil Nadu, India

²Department of Chemistry, Laboratory for Venom Peptidomics and Molecular Simulation, Bannari Amman Institute of Technology, Erode, Tamil Nadu, India

*Corresponding Author Email: arungowrishankar@gmail.com

Article Received on: 13/02/18 Approved for publication: 12/03/18

DOI: 10.7897/2230-8407.09340

ABSTRACT

Azadirachta indica (Neem) has a dynamic role in various problems associated with human health. Each part of the Neem tree has some medicinal property and is thus commercially utilizable. The plant parts such as leaves, flowers, seeds, fruits, roots and bark have been used traditionally for the treatment of inflammation, infections, fever, skin diseases and dental disorders. The aim of this study is to identify and characterize the vital bioactive compounds from the Neem sap by Gas chromatography and Mass spectroscopy (GC-MS). The GC-MS analysis of the Neem sap revealed the presence of 30 volatile compounds. Among the 30 compounds, the most predominant compounds are fatty acids like Hexadecanoic acid and Pentadecanoic acid. Hence, this current attempt forms a basis for the biological characterization and importance of the compounds which could be exploited for future development of drugs.

KEYWORDS: GC-MS analysis; Volatile compounds; *Azadirachta indica*.

INTRODUCTION

Azadirachta indica is a member of mahogany family, Meliaceae. The taxonomic classification of Neem is as follows: Kingdom: Plantae, Order: Rutales, Suborder: Rutinae, Family: Meliaceae, Subfamily: Melioideae, Tribe: Melieae, Genus: *Azadirachta*, Species: *indica*¹. Neem tree is mainly cultivated in the Indian subcontinent. It is consumed as a vegetable in some parts of the Asian subcontinent but mostly used as traditional medicine for centuries to cure multiple human diseases. For example, Neem components have been shown to have antifungal, anthelmintic, antibacterial, antiviral, anti-diabetic, contraceptive, and sedative effects^{2,3}. Additionally, its constituents are applied in alternative (Ayurveda, Unani, Homeopathy) and modern medicine, e.g., for the treatment of diverse infectious, metabolic, or cancer diseases^{4,5}. Many compounds have been found in the exudates of the, *Azadirachta indica* plant that have been used medically by humans. Studies of extracts from all major parts of Neem plant including leaves, flowers, fruits, and seeds, have shown promising chemopreventive and therapeutic effects in pre-clinical research⁶. So, this is now being widely used in pharmaceutical and cosmetics industries. Many compounds have been isolated from the extracts of various parts of the Neem tree. Quercetin and β -sitosterol, were the first polyphenolic flavonoids purified from the fresh leaves of neem and were known to have antibacterial and antifungal properties. Neem plays role as free radical scavenging properties due to rich source of antioxidant. Azadirachtin and Nimbolide showed concentration-dependent antiradical scavenging activity⁷.

The extract of *A. indica* leaves showed significant anti-inflammatory activity in cotton pellet granuloma assay in rats⁸. The leaf extract has shown virucidal activity⁹. Neem leaves also have the wound healing activity in both excision and incision wound models¹⁰. *Azadirachta indica* fruit skin extract and isolated ingredient Azadiradione showed significant antinociceptive property¹¹. The medicinal utilities have been described, especially for leaf, fruit and bark. But research on the plant part, sap has not been investigated yet. Hence, this study is undertaken to find out the volatile compounds present in the sap of *A.indica* by using Gas chromatography and Mass spectroscopy technique.

MATERIALS AND METHODS

COLLECTION OF NEEM SAP

The green fruits of Neem were collected from the trees of Sathyamangalam, Tamil Nadu. Then, the white fluid was taken out from the fruits and stored in the separate beaker.

GC-MS ANALYSIS

GC-MS analysis of the sap of *A.indica* was performed using Thermo GC- Trace ultra-version 5.0 and Thermo MS DSQ II. The equipment has a DB 35 – MS Capillary Standard non-polar column with dimensions of 30 mm \times 0.25 mm ID \times 0.25 μ m film. For GC-MS detection, an electron ionization system was operated in electron impact mode with ionization energy of 70 eV. Helium gas

was used as a carrier gas at a constant flow rate of 1 ml/min, and an injection volume of 1 µl was employed. The injector was operated at 250 °C and the oven temperature was programmed from 70 °C with an increase of 6 °C /min to 260 °C. Mass spectra were taken at 70 eV and the total GC-MS running time was 40.53 min. The peaks corresponding to the components were identified based on Willey and NIST libraries as well as comparison of their retention indices. The relative percentage amount of each component was calculated by comparing its average peak area to the total areas. The results obtained have been tabulated.

DISCUSSION

The present study was undertaken to find out the volatile bioactive compounds present in the sap of *A.indica* by using Gas chromatography and Mass spectroscopy. In this study, the Neem fruits were collected and the white colored fluid was separated out. The volatile compounds present in it was characterized by using GC-MS.

GC-MS ANALYSIS

The Neem sap was analysed for the detection of bioactive compounds in it by using GC-MS. The GC-MS profile of the sap of *A.indica* is represented in figure 2 which showed the prominent peaks for the presence of 30 bioactive compounds. The probable volatile compounds corresponding to the peaks with their retention time (RT), molecular formula, molecular weight (MW), structure and concentration (peak area %) are tabulated in Table 1.

From the analysis, it is revealed that there are 30 probable volatile compounds present in the Neem sap. These constituents belong to

15 classes of compounds including hydrocarbons, fatty acids, esters, alcohols, isothiocyanate, thiocyanate, pyrazine, aromatics, alkamides, cyanides, steroids, halocompounds, urea and N-hydroxyimine derivatives, unsaturated alkenamides, alkyne and indol.

From the above characterized compounds, Hexadecanoic acid has anti-tumor activity to human leukemic cells as well as murine cells¹². Hexadecanoic acid (palmitic acid) has been reported to induce NF-κB activation in HaCaT keratinocytes, which is an important pathway involved in cancer development. Besides, fatty acids exhibit cytotoxicity against HeLa cells, n-Hexadecanoic acid showed significant cytotoxicity against oral cancer (KB), breast cancer (MCF-7) and small cell lung cancer (NCI-H187)¹³. It also acts against inflammation¹⁴. Docosane owns antimicrobial activity against both gram positive and gram negative bacteria¹⁵. Docosenamide in its biosurfactant owes to its anticancerous and antiviral activities¹⁶.

CONCLUSION

The presence of various volatile bioactive compounds in the Neem sap justifies the use of this in treating various ailments. Hence, this result also enhances the traditional usage of *A.indica* which possesses a number of bioactive compounds. Therefore, it is recommended as a source of pharmaceutical importance and can be used as target for many known and unknown ailments. Future work on this study is to isolate and characterize the useful bioactive compounds for treating various diseases.































Figure 1: GC-MS Profile of *Azadirachta indica* sap

S.No	Retention Time	Compound name	Molecular Formula	Molecular weight	Area%	Structure
1	6.10	Hexadecanoic acid, 2,3-dihydroxypropyl ester	C19H38O4	330	0.13	
2	6.83	Docosane	C22H46	310	0.25	
3	7.90	1-Phenylpent-1-en-3-yne	C11H10	142	0.26	
4	9.67	Tetradecene	C14H28	196	0.25	
5	16.74	7-Methoxy-1-tetralone Ethylene Dithioketal	C13H16OS2	252	0.66	
6	18.47	2-(2-Thienylmethyleneamino)thiazole	C8H6N2S2	194	0.16	
7	19.85	2-Pentadecanone, 6,10,14-trimethyl- (CAS	C18H36O	268	0.17	
8	21.66	Pentadecanoic acid, 14-methyl-, methyl ester	C17H34O2	270	0.18	
9	22.84	3-Methoxy-9-phenylphenanthrene	C21H16O	284	0.23	
10	23.82	7,9-Di-tert-butyl-1-oxaspiro(4,5)deca-6,9-diene-2,8-dione	C17H24O3	276	0.22	

11	25.51	2-tert-Butyl-1,3-diphenyl-4,5,6,7-tetrahydroindole	C ₂₄ H ₂₇ N	329	0.33	
12	26.26	3-(4-Chlorobenzoyl)-6-methoxy-9-N-methyl-4-[2'-(4-chlorobenzoyl)ethyl]carbazole	C ₃₀ H ₂₃ Cl ₂ N ₃ O ₃	515	0.24	
13	27.47	Docosane	C ₂₂ H ₄₆	310	1.76	
14	28.73	Triacontane	C ₃₀ H ₆₂	422	0.18	
15	30.22	Hexanedioic acid, bis(2-ethylhexyl) ester	C ₂₂ H ₄₂ O ₄	370	2.37	
16	31.15	Nonacosane	C ₂₉ H ₆₀	408	0.18	
17	31.46	Dodecachloro-9,10-dihydrophenanthrene	C ₁₄ Cl ₁₂	588	0.47	
18	31.81	[1Z,2(1R,2R)]-1-(Trimethylstannyl)-2-vinylcyclopentyl)ethenyl 2,2,4,4-tetramethyl-1,3-dioxazolidine-3-carboxylate	C ₂₀ H ₃₅ NO ₃ Sn	457	0.39	
19	32.26	1,2-Benzenedicarboxylic acid, bis(2-ethylhexyl) ester (CAS)	C ₂₄ H ₃₈ O ₄	390	0.17	
20	33.13	(2RS)-1,3,8-trimethyl-4-propyl-5-ethyl-2-(1-hydroxyethyl)-7methoxycarbonyl-6, gamma-methylenecarbonyl-porphine	C ₃₆ H ₄₂ N ₄ O ₄	594	71.27	

21	34.17	6-(3,4-Difluorophenyl)-3-cyano-4-[N-[(4-chlorophenyl)(phenyl)methyl]piperazino]-2H-pyran-2-one	C ₂₉ H ₂₂ ClF ₂ N ₃ O ₂	517	0.58	
22	34.74	4-Amino-3-diethylamino-6-nitrobenzoic acid	C ₁₁ H ₁₅ N ₃ O ₄	253	0.71	
23	35.25	2,4-Difluoro-1-hydroxynaphthalene	C ₁₀ H ₆ F ₂ O	180	0.22	
24	35.78	1,4-Bis(bromomethyl)-2,6-dimethylbenzene	C ₁₀ H ₁₂ Br ₂	290	0.83	
25	36.11	m-Bis(3,4-diamino-2,5-diphenyl-6-pyridino)benzene	C ₄₀ H ₃₂ N ₆	596	0.37	
26	37.17	2,9-bis(2',6'-dimethoxyphenyl)-1,10-phenanthroline	C ₂₈ H ₂₄ N ₂ O ₄	452	0.76	
27	38.33	Nonane, 4-methylene	C ₁₀ H ₂₀	140	6.95	
28	39	13-Docosamide	C ₂₂ H ₄₃ NO	337	5.57	
29	39.92	1-[4-[[[(1,1,2-Trimethylpropyl)dimethylsilyl]oxy]methyl]phenyl]-2,2,2-trifluoro-1-ethanone O-tosyl oxime	C ₂₄ H ₃₂ F ₃ NO ₄ Si	515	0.13	


30	40.28	1-Trimethylsilyloxy-2-trimethylsilylamino-3-(3'-methoxy-4'-trimethylsilyloxy-phenyl) propanone	C19H37NO4S i3	427	3.99	
----	-------	--	---------------	-----	------	---

Table 1: Volatile components identified in the sap of *Azadirachta indica*

ABBREVIATIONS

GC-MS - Gas Chromatography Mass Spectrometry
 RT - Retention Time
 MW - Molecular Weight
 NIST - National Institute of Standards and Technology

REFERENCES

- Bhat S S. Neem – A Green Treasure. Electronic Journal Biology. 2008; 4(3):102–11.
- Veitch G E, Boyer A, Ley S V. The azadirachtin story. Angewandte Chemie - International Edition. 2008; 47: 9402–29.
- Puri H S. Neem The Divine Tree *Azadirachta indica*. Harwood Academic Publishers. 1999.
- Brahmachari G. Neem - An omnipotent plant: A retrospection. ChemBioChem. 2004; 5: 408–21.
- Lokanadhan S, Muthukrishnan P, Jeyaraman S. Neem products and their agricultural applications. Journal of Biopesticides. 2012; 5(SUPPL.):72–6.
- Paul R, Prasad M, Sah N K. Anticancer biology of *Azadirachta indica* L (neem): A mini review. Cancer Biology and Therapy. 2011; 12: 467–76.
- Hossain M A, Al-Toubi W A S, Weli A M, Al-Riyami Q A, Al-Sabahi J N. Identification and characterization of chemical compounds in different crude extracts from leaves of Omani neem. Journal of Taibah University for Science [Internet]. 2013; 7(4):181–8. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S1658365513000423>
- Chattopadhyay R R. Possible biochemical mode of anti-inflammatory action of *Azadirachta indica* A. Juss. in rats. Indian Journal of Experimental Biology. 1998; 36(4): 418–20.
- Parida M M, Upadhyay C, Pandya G, Jana A M. Inhibitory potential of neem (*Azadirachta indica* Juss) leaves on Dengue virus type-2 replication. Journal of Ethnopharmacology. 2002; 79(2): 273–8.
- Schumacher M, Cerella C, Reuter S, Dicato M, Diederich M. Anti-inflammatory, pro-apoptotic, and anti-proliferative effects of a

methanolic neem (*Azadirachta indica*) leaf extract are mediated via modulation of the nuclear factor- κ B pathway. Genes and Nutrition. 2011; 6(2):149–60.

- Ilango K, Maharajan G, Narasimhan S. Anti-nociceptive and anti-inflammatory activities of *Azadirachta indica* fruit skin extract and its isolated constituent azadiradione. Natural Product Research. 2013; 27(16): 1463–7.
- Harada H, Yamashita U, Kurihara H, Fukushi E, Kawabata J, Kamei Y. Antitumor activity of palmitic acid found as a selective cytotoxic substance in a marine red alga. Anticancer Research. 2002; 22(5): 2587–90.
- Keawsa-ard S, Liawruangrath B, Liawruangrath S, Teerawutgulrag A, Pyne S G. Chemical constituents and antioxidant and biological activities of the essential oil from leaves of *Solanum spirale*. Natural Product Communications. 2012; 7(7):955–8.
- Aparna V, Dileep K V., Mandal PK, Karthe P, Sadasivan C, Haridas M. Anti-Inflammatory Property of n-Hexadecanoic Acid: Structural Evidence and Kinetic Assessment. Chemical Biology and Drug Design. 2012; 80(3):434–9.
- Karabay-Yavasoglu N U, Sukatar A, Ozdemir G, Horzum Z. Antimicrobial activity of volatile components and various extracts of the red alga *Jania rubens*. Journal of Physical Therapy. 2007; 21(2):153–6.
- Donio MBS, Ronica S F A, Viji V T, Velmurugan S, Jenifer J A, Michaelbabu M, et al. Isolation and characterization of halophilic *Bacillus* sp. BS3 able to produce pharmacologically important biosurfactants. Asian Pacific Journal of Tropical Medicine. 2013; 6 (11):876–83.

Cite this article as:

Praveen Kumar P et al. Analysis of volatile compounds in the sap of *Azadirachta indica* (neem) using gas chromatography mass spectrometry. Int. Res. J. Pharm. 2018;9(3):38-43 <http://dx.doi.org/10.7897/2230-8407.09340>

Source of support: Nil, Conflict of interest: None Declared

Disclaimer: IRJP is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IRJP cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of IRJP editor or editorial board members.