



Research Article

SYNTHESIS AND THEORETICAL STUDY OF NEW QUINAZOLINONE DERIVATIVES CONTAINING VARIOUS HETEROCYCLIC MOIETIES

Hadel A. Abdulgane, Barakat A. F. Kamel *, Firyal W. Askar and Hamed H. Mohammed

Department of Chemistry, College of Science, University of Al-Mustansiriyah, Baghdad, Iraq

*Corresponding Author Email: baraka19832012@yahoo.com

Article Received on: 17/05/18 Approved for publication: 30/05/18

DOI: 10.7897/2230-8407.09576

ABSTRACT

In this work, quinazolinone derivatives were prepared, characterized by IR, UV, HNMR spectroscopy. Theoretically these compounds were studied using Hartree-Fock method and 3-21G basis set, where it was found HOMO-LUMO and thermodynamic functions (ΔH_f° , ΔG_f° and ΔS_f°) calculations, which found from values of these functions that the reactions of quinazolinone derivatives synthesis were exothermic reaction.

KEYWORDS: Quinazolinone Derivatives, IR, UV Spectroscopy, theoretical study and Hartree-Fock method

INTRODUCTION

Quinazolinone is an interesting and important aromatic heterocyclic system¹. Quinazolinone have been frequently used in medicine because of their wide range of biological activities². They are considered to be a potent structure for drug Developments³ Different quinazolinone derivatives have been reported for their antibacterial, antifungal⁴, anticancer⁵, anti HIV, anti-inflammatory, CNS depressants⁶ and anti tubercular activities^{7,8}.

MATERIALS AND METHODS

All of the reagents are commercially available in the (Aldrich Co.). Melting point was recorded by electric thermal melting point apparatus. FT-IR measurements have been recorded through the model Shimadzu FT-IR-8400S., UV. Spectra was measured by CARY 100, VARIAN Co. Model and H¹ NMR spectra using Ultra Shield Bruker laboratory model in the 300 MHz using DMSO-D₆ as a solvent and used of TMS, according to internal standards.

Synthesis of 2-Mercapto -3-(pyridine-2-yl)quinazolin-4(3H)-one (1)

Carbon disulfide (12ml) was added drop wise to an ice cold solution of potassium hydroxide (0.015 mol) in absolute ethanol (25 ml) containing solution of anthranilic (0.01mol) and (0.01mol) 2-amino pyridine .The mixture refluxed for 8 hours then poured into ice-water. The solid was filtered washed with water and re crystallized from ethanol. Yield: 78%. M.p. 180-182°C.

Synthesis of 2- Hydrazinenyl-3-(pyridine-2-yl)quinazolin-4(3H)-one (2)

A mixture of ethyl ester (0.01mol) and hydrazine hydrate 99 % (0.01mol) in absolute ethanol was heated under reflux for 8hrs

The excess of ethanol was distilled-off to a possible extent and cooled. The resultant solid was separated by filtration and dried. It was purified by re crystallization from ethanol. 85%. M.p. 199-201°C.

Synthesis of 2-(2-(4-bromo benzylidene)hydrazinyl)-3-(pyridine-2-yl)quinazolin-4(3H)-one (3)

A mixture of hydrazides (2) (0.0043mol), (0.0043mole) of the 4-Bromobezaldehyde and 0.1 ml of glacial acetic acid in (50ml) methanol was refluxed for 8 hrs. The formed precipitate was filtered, dried and purification from chloroform Yield: 73%. M.p.: 210--212°C.

Synthesis of 2-((2-(4-bromo phenyl)-3-chloro-4-oxoazetidin-1-yl)amino)-3-(pyridine-2-yl)quinazolin-4(3H)-one (4)

A mixture of Schiff base(3) (0.01 mol) and tri ethyl amine (0.02 mol) was dissolved in Dioxane (20 ml). chloro acetyl chloride (0.02 mol) was added in portion wise with vigorous shaking at room temperature for 30 min. The reaction mixture was heated under reflux for 5h and the content was kept at room temperature for 24 h and poured into ice-cold water. The product obtained was filtered, washed several times with water. Yield: 78%. M.p.: 226-228°C.

Synthesis of 2-(4-bromo phenyl)-3-((4-oxo-3-(pyridine-2-yl)-3,4-dihydroquinazolinyl)amino)thiazolidin-4-one (5)

A mixture of Compound (3) (0.001mol) was solved in 30mL chloroform with ZnCl₂ (0.01g) and (0.005mol) of thio glycolic acid was added to the mixture, the mixture was refluxed for 10h. The reaction completion was monitored by thin layer chromatography (TLC) using ethyl acetate : hexane system (3:7). The solvent was removed under reduced pressure, residue treated by solution of 10% NaHCO₃ to removed excess of mercapto

acetic acid, washed with water, dried and re crystallization from acetone. Yield: 70%.M.p. 182-184°C.


Antimicrobial Activity

The antimicrobial activity of the quinazolinone derivatives were tested by the agar disc-diffusion method against two Gram positive *Staphylococcus aureus* and *Streptococcus pyogenes* and two gram negative bacteria *E. Coli*, *Klebsiella* spp micro organisms and fungal strains namely *Candida albicans*. Dimethyl sulfoxide DMSO was used as a control and as solvent .The test was performed at 100mg/mL concentration. The bacteria and fungi were carried out in agar and potato dextrose


agar medium and these plates were incubated for 24 h for bacteria and 48h for fungi at 37 °C.

Computational Details


Hartree –Fock method were performed with the Gaussian 09 series of programs. The total energy of the prepared compounds was calculated at Basis Set 3-21G. by this method (IR, UV) spectrum and thermodynamic functions (ΔH°_f , ΔG°_f and ΔS°_f) were studied⁹. Figure1 shows the HOMO and LUMO for the compounds:




Compound 1




Compound 2



Compound 3



Compound 4




Compound 5

Figure 1: HOMO-LUMO Categories of the prepared compounds

RESULTS AND DISCUSSION

The new quinazolinone derivatives were prepared following the reaction sequences depicted in Scheme 1.




Scheme 1

FTIR Spectra

The 2-hydrazineyl-3-(pyridin-2-yl)quinazolin-4(3H)-one (2) was prepared by refluxing of compound 1 with hydrazine hydrate in absolute ethanol for 8 h. The FTIR spectrum indicated the appearance two bands in the region (3363-3176) cm⁻¹ which could be attributed to asymmetric and symmetric stretching vibration of NH₂ and NH. Condensation of the compound (2) with *p*-bromobenzaldehyde in absolute ethanol afforded the Schiff base (3). The formation of Schiff base was indicated by the presence in their FTIR spectra of azomethine (CH=N) stretching band at (1616) cm⁻¹, combined with the disappearance of the NH₂ stretching band. Furthermore, the generation of azetidinyl derivative (4) by the reaction of Schiff base (3) with

Et₃N and chloroacetyl chloride (C₂H₂OCl₂) in dioxane. The FT-IR spectrum indicated the appearance of band at 1728 cm⁻¹ (C=O) due to β-lactam with disappearance band of imine (N=CH) in the region 1616 cm⁻¹. Cyclization of Schiff base (3) with mercaptoacetic acid in the presence of chloroform as solvent afforded thiazolidenone (5). The structures of compound (5) was confirmed by the presence of carbonyl stretching band at (1705) cm⁻¹ due to thiazolidenone ring which was the characteristic evidence for success of cyclization. Figures 2, 3 and table 1 showed the FTIR spectrum of the compounds 3 and 5.






Figure3: FTIR Spectra of compound 5


Table1: Experimental and Theoretical values of FTIR Spectrum bands of prepared compounds

Compound	ν (C=O)	ν (CH=N)	ν (N-H)	ν (C=S)
Comp.1	1685 (1693) _H	-----	3367 (3381) _H	1610 (1607) _H
Comp.2	1656 (1670) _H	-----	3363-3176 (3367) _H	-----
Comp.3	1690 (1694) _H	1620 (1616) _H	3210 (3348) _H	-----
Comp.4	1687(amide) (1640) _H 1728(lactam) (1727) _H	-----	3340 (3321) _H	-----
Comp.5	1725 (1720) _H	-----	3328 (3304) _H	-----


H: Hartree-Fock

UV. Spectra

The UV spectra of the prepared compounds showed in figure 4 and table 2:



1



5

Figure4: UV Spectra of the compounds 1 and 5

Table2: Experimental and Theoretical bands of UV Spectrum of prepared compounds

Compound	Band I	Band II
Comp. 1	241 (243) _H	209 (211) _H
Comp.2	247 (245) _H	206 (215) _H
Comp.3	292 (301) _H	225 (223) _H
Comp.4	290 (309) _H	232 (227) _H
Comp.5	293 (303) _H	227 (218) _H

H: Hartree-Fock

¹H NMR Spectra

¹H NMR spectrum of compound 1 (δ , ppm): 9.23(s, 1H, NH), 8.46-6.45 (m, 8H, Ar-H). compound (2) exhibited singlet signal at 3.5ppm and at 9.38 ppm was assigned to NH proton, signal at 5.71 ppm was attributed protons of NH₂, aromatic protons were appeared at 8.0-7.05ppm. The ¹H NMR of derivative (3) showed singlet signal at 8.7ppm due to NH proton, while aromatic protons and proton of CH=N appears as

multiple signals at 8.1-6.2 ppm. For compound 4, ¹H NMR (300 MHz, δ , ppm, DMSO-d₆): 9.45 (s, 1H, NH), 8.45-6.52 (m, 8H, Ar-H), 5.5-5.0 (dd, 2H, CH). Compound 5 appears 8.78 (s, 1H, NH), 8.53-6.46 (m, 8H, Ar-H), 5.40 (s, 1H, CH-N), 3.83 (s, 2H, CH₂). Figures 5 and 6 show the ¹H NMR spectrum of compounds 2 and 3.




Figure 5: ¹H NMR Spectra of compound 2




Figure 6: ¹H NMR Spectra of compound 3

Antimicrobial Activity

According to table 3 compound 1 showed that it had no effect of inhibition on *Staphylococcus aureus*, while compound 2 showed that it had no effect of inhibition on *Klebsiella spp* microorganisms. Compounds 3, 4 and 5 showed inhibition for all kinds of bacteria higher than other compounds.

Table3: Antimicrobial Evaluation of prepared compounds

Heterocyclic Derivative	inhibition zone (mm) at 100 mg/mL				
	Gram positive		Gram negative		Fungi
	<i>S. aureus</i>	<i>S. epidermidis</i>	<i>E. coli</i>	<i>Klebsiella spp</i>	<i>C. albicans</i>
Comp.1	-	14	16	15	15
Comp.2	15	12	16	-	15
Comp.3	16	15	14	11	18
Comp.4	18	13	12	20	15
Comp.5	20	19	17	22	19

Thermodynamic Properties

Thermodynamic properties of the prepared compounds were calculated theoretically using Hartree-Fock method by Gaussian 09 program. Table 4 shows the thermodynamic functions (ΔH°_f , ΔG°_f and ΔS°_f) of these compounds:

Table4: Thermodynamic properties of prepared compounds

Comp.	ΔH°_f (kcal/mol)	ΔG°_f (kcal/mol)	ΔS°_f (cal/mol.K ⁻¹)
Comp.1	-223546.494	-223528.924	-58.959
Comp.2	-840.443	-840.491	-0.161
Comp.3	-3665.912	-3665.974	-0.208
Comp.4	-4268.994	-4269.051	-0.191
Comp.5	-4211.036	-4211.101	-0.218

Above table showed that the reactions of compounds synthesis were exothermic reactions.

CONCLUSIONS

we concluded from the comparison between the experimental and theoretical results using HF method with basis set 3-21G that there is a simple difference between them the reason is that these programs operate in standard conditions and in a gas situation. From the study of antimicrobial activity of quinazolinone derivatives on the different kinds of bacteria showed that these compounds had significant effect in the inhibition of these bacteria. The values of thermodynamic function (negative enthalpy, negative entropy and negative free energy) showed that the synthesis reactions of quinazolinone derivatives from compound 1 to compound 5 were (exothermic reactions).

ACKNOWLEDGEMENT

All the experiments of this work were carried out in the laboratories of department of chemistry in the university of al-mustansiriyah.

REFERENCES

- 1.Firyal W. Askar, Salma A. R. Abbas, Redha I. Al-Bayati and Hanaa A. Al-Tamemi. Synthesis and biological evaluation of new quinazolinone derivatives. European Journal of Chemistry 2014; 5 (4): 628-634.
- 2.Sudhakar Rao G., Kalaichelvan VK. and Sudhakar Rao G. Synthesis and Antibacterial Activity of 3-phenyl substituted Quinazolinone Derivatives Via Chalcones. IJRPC 2015; 5(3): 470-474
- 3.Vijayakumar K., Jafar Ahamed A. and Thiruneelakandan G.. Synthesis, Antimicrobial, and Anti-HIV1 Activity of Quinazoline-4(3H)-one Derivatives. Journal of Applied Chemistry 2013; 2013.
- 4.Indu S., and Arun K. Synthesis and Antimicrobial activity of various Quinazolinone derivatives containing Thiazole and Thiazolidinone Moiety. International Journal of ChemTech Research. 2014; 6(5): 2649-2655.
- 5.Ghadamali K., Elham J., Gholamhossein H., Daryoush A., Marzieh R. K. and Farshid H., Synthesis of Some New Quinazolinone Derivatives and Evaluation of Their Antimicrobial Activities, Iranian Journ Pharmaceutical Research 2012; 11 (3): 789-797.
- 6.Rajveer CH., Kumaraswamy D., Sudharshini S. and Stephen R. B. Synthesis of some 6- bromo quinazolinone Derivatives for Their Pharmacological Activities. International Journal of Pharma and Bio Sciences. 2010; 1(3).
- 7.Maneesh K. S. and Shantakumar S. M. Design and Synthesis of Novel 2-Trichloromethyl-4Substituted Quinazoline Derivatives as Anti-tubercular Agents. Chem Sci Trans 2013; 2(3): 1056-1062.
- 8.Mohammad A. Chemical Characteristics, Synthetic Methods, and Biological Potential of Quinazoline and Quinazolinone Derivatives. International Journal of Medicinal Chemistry 2014; 2014.
- 9.María V. R., Manuel T., Pilar J., Rafael N., James S. C., Ana F. L. O. M. S., Manuel A. V. R. Thermo chemistry of 2- and 3-Acetylthiophenes: Calorimetric and Computational Study. J. Phys. Chem. 2007; 111 (43): 11084–11092.

Cite this article as:

Hadel A. Abdulgane *et al.* Synthesis and theoretical study of new quinazolinone derivatives containing various heterocyclic moieties. Int. Res. J. Pharm. 2018;9(5):75-80 <http://dx.doi.org/10.7897/2230-8407.09576>

Source of support: Nil, Conflict of interest: None Declared

Disclaimer: IRJP is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IRJP cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of IRJP editor or editorial board members.