

## SPECTROPHOTOMETRIC DETERMINATION OF CEPHALEXIN USING NINHYDRIN REAGENT IN TABLET DOSAGE FORM

Patel Satish A\*, Patel Natavarlal J.

Shree S. K. Patel College of Pharmaceutical Education &amp; Research, Ganpat University, Kherva, Mehsana, Gujarat, India

Article Received on: 13/07/11 Revised on: 18/08/11 Approved for publication: 10/09/11

\*Satish A. Patel, Department of Pharmaceutical Quality Assurance, Shree S. K. Patel College of Pharmaceutical Education & Research, Ganpat University, Kherva – 382711, Mehsana, Gujarat, India. Email: satishpatel\_77@yahoo.com

### ABSTRACT

A simple, sensitive, accurate, precise and economical visible spectrophotometric method was developed and validated for the estimation of cephalexin in tablets. The method is based on the reaction of cephalexin with ninhydrin reagent in methanol giving blue color chromogen, which shows maximum absorbance at 576 nm against reagent blank. The chromogen obeyed Beer's law in the concentration range of 5-60 µg/ml for cephalexin. The results of the analysis have been validated statistically and by recovery studies.

**Key words:** Cephalexin, Chromogen, Ninhydrin reagent, Visible spectrophotometric, Tablet.

### INTRODUCTION

Cephalexin (CFD) is chemically, (6*R*,7*R*)-7-[[*(2R)*-Aminophenylacetyl]amino]-3-methyl-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid<sup>1</sup>. Cephalexin is a first-generation cephalosporin antibacterial for the treatment of susceptible infections including those of respiratory tract, urinary tract and skin<sup>2</sup>. Cephalexin is official in IP, USP and BP. IP<sup>3</sup>, USP<sup>4</sup> and BP<sup>5</sup> describes liquid chromatography method for the estimation of cephalexin. Literature survey reveals polarography<sup>6</sup>, spectrophotometric<sup>7-9</sup>, fluorimetric<sup>10</sup>, HPTLC<sup>11</sup> and HPLC<sup>12</sup> methods for the estimation of cephalexin in biological fluids and in pharmaceutical formulations. The present communication describes simple, sensitive, accurate, precise and economical visible spectrophotometric method using ninhydrin reagent for the estimation of cephalexin in tablet dosage form.

### MATERIALS AND METHODS

#### Apparatus

A Shimadzu model 1601 double beam UV/Vis. spectrophotometer with spectral width of 2 nm, wavelength accuracy of 0.5 nm and a pair of 10 mm matched quartz cells was used to measure absorbance of the resulting solutions. A Sartorius CP224S analytical balance (Germany), an ultra sonic cleaner (Frontline FS 4, Mumbai, India), water bath (Cintex Industrial Corporation, Mumbai, India) were used in the study.

#### Reagents and Materials

Cephalexin powder was procured as a gift sample from Mann Pharmaceuticals Ltd, Mehsana, Gujarat, India. The commercially available tablets of cephalexin were procured from local market. Ninhydrin reagent and methanol (AR Grade, S.D. Fine Chemicals Ltd., Mumbai) were used in the study.

#### Preparation of reagent and standard stock solution

Accurately weighed ninhydrin powder (500 mg) was transferred to a 100 ml volumetric flask, dissolved in and diluted to the mark with methanol (0.5% w/v). Accurately weighed cephalexin (20 mg) was transferred to a 100 ml volumetric flask, dissolved in and diluted to the mark with methanol. (200 µg/ml).

#### Methodology

Standard stock solution of cephalexin (1.0 ml) was transferred to a 10 ml corning volumetric flask. Ninhydrin reagent (2.0 ml) was added and mixed. The flask was immersed in a water bath at 92 ± 1<sup>0</sup> C for 20 minutes, cooled to room temperature and the volume was adjusted to 10 ml with methanol. The absorbance of the colored solution was scanned in the range of 400 to 800 nm against reagent blank, prepared similarly in which volume of standard solution was

replaced by an equal volume of methanol. Maximum absorbance was obtained at 576 nm.

#### Optimization of different conditions

##### Effect of concentration of ninhydrin reagent

Standard stock solution of cephalexin (1.0 ml) was transferred to a series of 10 ml corning volumetric flasks. To each flask, different volumes of ninhydrin reagent (0.5, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5 and 4.0 ml) was added and mixed. The flasks were immersed in a water bath at 92 ± 1<sup>0</sup> C for 20 minutes, cooled to room temperature and the volume in each flask was adjusted to 10 ml with methanol. The absorbance of the resulting solutions was measured at 576 nm against reagent blank. Maximum absorbance was observed in the presence of 2.0 ml of 0.5% w/v ninhydrin reagent.

##### Effect of temperature

Standard stock solution of cephalexin (1.0 ml) was transferred to a series of 10 ml corning volumetric flasks. To each flask, 2.0 ml of ninhydrin reagent was added and mixed and heated at different temperatures (40<sup>0</sup>, 50<sup>0</sup>, 60<sup>0</sup>, 70<sup>0</sup>, 80<sup>0</sup>, 90<sup>0</sup>, 92<sup>0</sup>, 94<sup>0</sup> and 96<sup>0</sup> C) for 20 minutes, cooled to room temperature and the volume in each flask was adjusted to 10 ml with methanol. The absorbance of the resulting solutions was measured at 576 nm against reagent blank. Optimum temperature to obtain maximum absorbance was found to be 92 ± 1<sup>0</sup> C.

##### Time for maximum color development and color stability

Standard stock solution of cephalexin (1.0 ml) was transferred to a series of 10 ml corning volumetric flasks. To each flask, 2.0 ml of ninhydrin reagent was added and mixed. The flasks were immersed in a water bath at 92 ± 1<sup>0</sup> C for different time interval (5, 10, 15, 20, 25, 30, 45 and 60 minutes), cooled to room temperature and the volume in each flask was adjusted to 10 ml with methanol. The absorbance of the resulting solutions was measured at 576 nm against reagent blank. Maximum absorbance was obtained after 20 minutes, which remain constant for 1 hour.

##### Validation of the proposed method

The proposed method was validated according to the International Conference on Harmonization (ICH) guidelines<sup>13</sup>.

##### Linearity

Calibration curve was plotted over a concentration range of 5-60 µg/ml for cephalexin. Accurately measured standard stock solutions of cephalexin (0.25, 0.5, 1.0, 1.5, 2.0 and 3.0 ml) were transferred to a series of 10 ml corning volumetric flasks. To each flask, 2.0 ml of ninhydrin reagent was added and mixed. The flasks were immersed in a water bath at 92 ± 1<sup>0</sup> C for 20 minutes, cooled to room temperature and the volume in each flask was adjusted to 10 ml with methanol. The absorbance of the resulting solutions was measured at

576 nm against reagent blank. Calibration curve was constructed for cephalixin by plotting concentration versus absorbance at 576 nm. Each reading was an average of five determinations.

#### Accuracy (% Recovery)

The accuracy of the method was performed by calculating recovery of cephalixin by the standard addition method. Known amounts of standard solutions of cephalixin were added at 50, 100 and 150% levels to prequantified sample solutions of cephalixin (20 µg/ml). Each sample was prepared in triplicate at each level. The amount of cephalixin was estimated by applying obtained values to regression equation.

#### Method Precision (% Repeatability)

The precision of the instrument was checked by repeated scanning and measurement of the absorbance of solutions (n = 6) of cephalixin (20 µg/ml) without changing the parameters for the method. The repeatability was expressed in terms of relative standard deviation (%RSD).

#### Intermediate Precision (Reproducibility)

The intraday and interday precision of the proposed method was performed by analyzing the corresponding responses 3 times on the same day and on 3 different days over a period of 1 week for different concentrations of standard solutions of cephalixin (5-60 µg/ml). The results were reported in terms of relative standard deviation (%RSD).

#### Limit of Detection (LOD) and Limit of Quantification (LOQ)

The limit of detection (LOD) and the limit of quantification (LOQ) of the drug was derived by calculating the signal-to-noise ratio (S/N, i.e., 3.3 for LOD and 10 for LOQ) using the following equations as per International Conference on Harmonization (ICH) guidelines<sup>14</sup>.

$$\text{LOD} = 3.3 \times \sigma/S$$

$$\text{LOQ} = 10 \times \sigma/S$$

Where  $\sigma$  = the standard deviation of the response and S = Slope of calibration curve.

#### Estimation of cephalixin from pharmaceutical tablet dosage form

Twenty tablets were accurately weighed and powdered. A quantity of powder equivalent to 20 mg of cephalixin was transferred to a 100 ml volumetric flask and mixed with methanol (50 ml) and sonicated for 20 minutes. The solution was filtered through Whatman filter paper No. 41 and the residue was washed thoroughly with methanol. The filtrate and washings were combined in a 100 ml volumetric flask and diluted to the mark with methanol. The solution (2.0 ml) was transferred to a 10 ml corning volumetric flask. Ninhydrin reagent (2.0 ml) was added and mixed. The flask was immersed in a water bath at  $92 \pm 1^{\circ}$  C for 15 minutes, cooled to room temperature and the volume was adjusted to 10 ml with methanol. The absorbance of the resulting solution was measured at 576 nm against reagent blank. The amount of cephalixin present in tablet solution was determined by fitting the responses into the regression equation. The analysis procedure was repeated five times with tablet formulation.

#### RESULTS AND DISCUSSION

Ninhydrin reagent is mainly used for the detection of free amino and carboxyl groups in proteins and peptides, yielding a blue color under the proper condition<sup>14</sup>. Ninhydrin (triketohydrinden hydrate) reacts with amino group containing substances like amino acids, proteins and peptides and when heated under proper conditions, produce ammonia, carbon dioxide and blue purple complex<sup>15-18</sup>. Literature survey reveals spectrophotometric method for the determination of lisinopril using ninhydrin reagent<sup>19</sup>. Therefore it was thought of interest to extend the application of ninhydrin reagent in the estimation of amino group containing drugs like cephalixin.

In the proposed method, reagent solution and standard stock solutions of drugs were prepared in methanol. Various reaction conditions were established by varying one parameter at a time and

keeping the others fixed by observing the effect produced on the absorbance of the colored species. The parameters involved for maximum color development viz. concentration of ninhydrin reagent, temperature and heating time required to yield chromogen of maximum color intensity and stability were optimized. In this method all these parameters were strictly followed.

The blue colored complex formed having wavelength of maximum absorbance at 576 nm (Figure 1). In proposed method, it was found that 2.0 ml of 0.5% w/v ninhydrin reagent (Figure 2),  $92 \pm 1^{\circ}$  C heating temperature (Figure 3) and 15 minutes heating time (Figure 4) was sufficient for the development of maximum color intensity. Stability study of the developed chromogen was carried out by measuring the absorbance values at a time intervals of 20 minutes for 3 h and it was found to be stable for more than 2 h for the drugs at room temperature.

The linearity was found in the concentration range of 5 to 60 µg/ml ( $r^2 = 0.9958$ ). The reproducibility, repeatability and precision of method are very good as shown by the low values of standard deviation and relative standard deviation (%RSD). The % recovery value in the range of 98.92 to 102.2 % for tablet indicates non-interferences from the formulation excipients. The data of recovery studies and assay results are given in Table 1 and Table 2, respectively. Optical characteristics of method and summary of validation parameters for cephalixin was given in Table 3.

#### CONCLUSION

The proposed visible spectrophotometric method was found to be, simple, sensitive, accurate, precise and economic for determination of cephalixin in tablet dosage form. Hence it can be conveniently adopted for routine quality analysis of the drug in pharmaceutical dosage form.

#### ACKNOWLEDGEMENT

The authors are thankful to Mann Pharmaceutical Limited, Mehsana, Gujarat, India for providing gift sample of cephalixin for research. The authors are highly thankful to Shree S. K. Patel College of Pharmaceutical Education & Research, Ganpat University, Kherva, Mehsana, Gujarat, India for providing all the facilities to carry out the work.

#### REFERENCES

- Maryadele. J. O' Neil. The Merck Index: An Encyclopedia of chemicals, drugs and biologicals, 14th ed. New Jersey: Published by Merck Research Laboratories, Division of Merck and Co., Inc. Whitehouse station: 2006. p. 326.
- Sweetman SC, Eds. In The Martindale: The Complete Drug Reference. 35th ed. London, UK: Pharmaceutical Press; 2007. p. 194.
- The Indian Pharmacopoeia, Vol. II, Government of India, Delhi: The Controller of Publications; 2010. p. 1030.
- British Pharmacopoeia, Vol. I, Stationary Office, London: Medicines and Healthcare Products Regulatory Agency, 2010. p. 397.
- The United States Pharmacopoeia, USP 32, NF 27, Vol. 2, Rockville, MD: The United States Pharmacopoeial Convention, Inc., 2009. p. 1870.
- Fogg AG, Fayad NM. Differential pulse polarographic study of the degradation of cephalixin: Determination of hydrogen sulphide and other degradation products. *Anal Chimica Acta* 1979;110:107-115.
- Alwarthan AA, Fattah SA, Zahran NM. Spectrophotometric determination of cephalixin in dosage forms with imidazole reagent. *Talanta* 1992;39:703-707.
- Patel SA, Patel NM, Patel MM. Spectrophotometric methods for the estimation of Cephalixin in tablet dosage forms. *Indian J Pharm Sci* 2006;68:278-280.
- Al-Ghannam SM. Spectrophotometric and atomic absorption spectrometric determination of cephalixin and cephradine in dosage forms. *J Food Drug Anal* 2008;16:19-25.
- Barbhajia RH, Turner P. Fluorimetric determination of ampicillin and cephalixin. *Br J Pharmacol* 1976;58:473-475.
- Coran SA, Bambagiotti-Alberti M, Giannellini V, Baldi A, Picchioni G, Paoli F. Development of a densitometric method for the determination of cephalixin as an alternative to the standard HPLC procedure. *J Pharm Biomed Anal* 1998;18:271-274.
- Pecavar A, Smidovnik A, Milivojevic D, Prosek, M. A reversed phase high-performance liquid chromatography method for determination of cephalixin in human plasma. *Journal of High Resolution Chromatography*, 1997;20:674-678.
- The International Conference on Harmonization, Q2 (R1), Validation of Analytical Procedure: Text and Methodology, 2005.
- Budavari S. The Merck Index, 13th ed. Merck & Co., Inc., Whitehouse Station, NJ: 2001. p. 1175.

- 15) Holme D.J. and Peck H. Analytical Biochemistry, 3rd ed. The Addition Wesley Longman Ltd, New York: 1998, p. 356.
- 16) Mathews C.K. and Van Holde K.E. Biochemistry, 2nd ed. The Benjamin/Cummings Publishing Co. Inc. Menlo Park, California: 1996, p. 156.
- 17) Metzler D.E. Biochemistry: The Chemical Reactions of Living Cells, 2nd ed. Volume 1, The Harcourt Academic Press, London: 2001, p. 120.
- 18) Berg J.M. Tymoczko J.L. and Stryer L. Biochemistry, 5th ed. The W. H. Freeman and Co. New York: 2002, p. 91.
- 19) Mashru R, Parikh P. Simple spectrophotometric analysis of lisinopril. Eastern Pharmacist 2001;528:113-115

TABLE 1: RESULTS OF RECOVERY STUDIES IN TABLET AND INJECTION DOSAGE FORMS

Formulation	Level	Amount of sample taken (µg/ml)	Amount of standard spiked (%)	Mean % recovery ± S. D.* (n = 3)
Tablet	I	20	50	98.92 ± 1.57
	II	20	100	102.2 ± 1.41
	III	20	150	100.7 ± 0.92

S. D. is standard deviation and n is the number of determinations.

TABLE 2: RESULTS OF ANALYSIS OF TABLET FORMULATION

Formulation	Label claim (mg)	Amount found (mg)	% Label Claim ± S. D.* (n = 5)
Tablet	Brand I	250	249.6
	Brand II	500	509.0
			99.84 ± 1.08
			101.8 ± 1.57

S. D. is standard deviation and n is the number of determinations.

TABLE 3: OPTICAL CHARACTERISTICS AND SUMMARY OF VALIDATION PARAMETERS

Parameters	Results
λ max (nm)	576
Linearity range (µg/ml)	5 - 60
Sandell's sensitivity (µg/cm <sup>2</sup> /0.001 A.U.)	0.0485
Molar extinction coefficient (l/mol.cm)	7156
Correlation coefficient (r <sup>2</sup> )	0.9958
Regression equation (y* = b + ac)	
Slope (a)	0.0186
Intercept (b)	0.0423
Standard deviation (S. D.)	± 0.004693
% Relative standard deviation (% RSD)	± 1.216
Standard error of mean (S.E.M)	± 0.002103
Repeatability (% RSD, n = 6).	1.12
Intermediate Precision (% RSD)	
Interday (n = 3)	0.39-1.62
Intraday (n = 3)	0.43-1.98
Accuracy (% Recovery) (n = 5)	98.92 - 102.2
Limit of detection (LOD) (µg/ml)	1.22
Limit of quantification (LOQ) (µg/ml)	4.03

y\* = b + ac where 'c' is the concentration and y is absorbance unit. n is the number of determinations, RSD is relative standard deviation and S.E.M is standard error of mean..




Figure 1: Spectra of cephalixin with ninhydrin reagent at 576 nm in methanol




Figure 2: Optimization of volume of 0.5% w/v ninhydrin reagent (ml)




Figure 3: Optimization of heating temperature (°C)




Figure 4: Optimization of heating time (minutes)

Source of support: Nil, Conflict of interest: None Declared