



## Research Article

### EVALUATION OF ANTI MICROBIAL ACTIVITY OF ETHANOL EXTRACT OF *PSEUDARTHRIA VISCIDA* LEAF

Kavitha K \*1, Sangeetha N1 Nepolean R, Hari Babu Y

Department of Pharmaceutical chemistry, Thanthai Rover College of Pharmacy, Perambalur, Tamilnadu, India

Department of Pharmaceutical Chemistry, Grace College of Pharmacy, Athalur, Kodunthirapully, Palakkad, Kerala, India

\*Corresponding Author Email: kavithakrocks@gmail.com

Article Received on: 07/06/18 Approved for publication: 29/06/18

DOI: 10.7897/2230-8407.096114

#### ABSTRACT

The plant *Pseudarthritis viscida* (Linn) belongs to Fabaceae family. It is a perennial shrub distributed throughout all districts of south India, also reported from Sri Lanka and Timor. Traditionally, the plant used in treatment of intermittent fever, urinary diseases, tumors, edema, and burning sensation, difficulty in breathing and toxic conditions. Herbal medicine is used frequently a part of larger therapeutic system like Traditional and folk medicine. Assortment of literature, the plant possess anti-diabetic, anti oxidant, anti tumor, anti hypertensive, anti diarrheal activities and it is necessary to evaluate in a scientific base, the potential use of folk medicine for the treatment of many diseases. The present study was performed to evaluate the in vitro antimicrobial activity of ethanol extract leaves of *Pseudarthritis viscida* Linn, Phytochemical screening which indicated presence of carbohydrates, alkaloids, phenols, flavonoids.

**Key words:** *Pseudarthritis viscida* leaves, ethanol extract, microbial activity, carbohydrates,

#### INTRODUCTION

Natures have been a source of medicinal agents for thousands of years, a stinking of modern drug has been isolated from natural sources, some based on their use in traditional medicines or Phytomedicines. Scientific interest in medicinal plants has flourished due to enhanced efficiency of new plant derived drugs, growing interest in natural products, rising concerned the side effects conventional medicine. The various use of herbal remedies, preparations are described throughout human history represents the origin of modern medicine. Many of conventional drugs originate from plant sources<sup>1</sup>.

The roots are astringent, digestive, constipated, anthelmintic, emollient, thermogenic, anti inflammatory, aphrodisiacs, cardiogenic, and febrifuge, also used for rejuvenating tonic. The whole plant exhibits anti oxidant, analgesic, anti fungal, anti pyretic effects. They are useful and vitiated the conditions of cough, asthma, bronchitis, tuberculosis, gout, hyperthermia, fever, general ability<sup>3</sup>. The root, leaf, stem, callus extract of the *Pseudarthritis viscida* exhibited significant anti fungal property<sup>4</sup>. The preliminary photochemical screening indicates that the presence of alkaloids, flavonoids, tannins and phenolic compounds, saponins. The ethanol extract of *Pseudarthritis* (Linn) root exhibited anti oxidant activity<sup>5</sup>. Phytochemical constituents like vaffeic acid, gallic acid, rutin and ferulic acid were identified from *Pseudarthritis viscida* roots by HPLC with UV detection at 280nm<sup>6</sup>. The methanol extract of *Pseudarthritis viscida* root exhibited 43 compounds such as cis vaccenic acid, gamma sitosterol and stigma sterol by GCMS. The ethanol extract of roots showed anti diabetic, anti inflammatory, diuretic effect.<sup>7</sup>, literature survey suggested that, to investigate and focus for the future scope of anti microbial activity on *Pseudarthritis viscida* leaf.



Fig 1: *Pseudarthritis viscida* leaf (Linn)

Kingdom: Plantae,  
Synonyms: Hedysarum viscidum  
Division: magnoliophytaclass, magnoliopsida.  
Genus: Pseudarthritis  
Species: viscida  
order: Fabale  
Common name: chappakno, nirmalli, moovila, salapami, neyakuponna.

*Pseudarthritis* is an under shrub, which is perennial, contain lupeol triterpenoid and it is semi erect in appearance., arranged alternately and contains three foliate leaves, also contains membranous leaflets, pink or rose coloured flowers<sup>2</sup>.

#### MATERIALS AND METHODS

The whole plant was collected from Kolli hills, Namakkal District, Tamil Nadu. The plant was authenticated by Botanical survey of India, Southern Circle, Coimbatore and a voucher specimen were deposited (No.BSI/SC//23/06-07/tech-166). The collected Leaves of *Pseudarthritis viscida* (Linn) was dried under shade drying, coarse powder was used for the extraction and phytochemical analysis.

### Phytochemical studies

Leaves of *Pseudarthritis viscida* Linn reduced to fine powder (# 42 size meshes) and stored in an air tight container and around 300 gm of powder were subjected to successive hot continuous extraction (Soxhlet) (60-80°C) with petroleum ether, ethanol(95%v/v). Finally, the drug will be air dried in hot air oven below 500°C. Later the effective extraction, the solvent was distilled off, then it was concentrated on water bath, the extracts obtained with each solvent have been weighed<sup>8</sup>. The various extract of leaves of *Pseudarthritis viscida* was subjected to chemical tests for identification of its active constituents.

### ANTIMICROBIAL STUDIES<sup>9,10</sup>

The 25,50,75 and 100µg/ml of ethanol extract were used to find out the anti microbial properties using gentamycin and fluconazole (10µg/ml) as a standard against various strains of gram positive, gram negative bacteria and fungi organism. The nutrient agar medium was prepared and sterilized by autoclave at 15 lbs pressure 120°C for 15 minutes, then aseptically poured the medium into the sterilized Petri plates and allowed to solidify the

bacterial, fungal broth culture was swabbed on each Petri plates using sterile buds. Then wells were made using cork borer on the solidified medium. The ethanol extract of plant leaves to each well aseptically. The same techniques were repeated for each Petri plates were incubated at 37°C for 24 hr after incubation the plates were observed zone of inhibition.

### RESULTS AND DISCUSSION

The present study on the ethanol extract leaves of *Pseudarthritis viscida* (Linn). The Phytochemical screening showed the presence of carbohydrates, alkaloids and flavonoids, amino acids and tannins, steroids. The concentration 100µg/ml of ethanol extract of *Pseudarthritis viscida* leaf exhibited significant activity against antibacterial and fungal organisms when compared to standard. The concentration of 50µg/ml & 75µg/ml showed moderate antibacterial, anti fungal activity when compared to standard. The concentration 25µg/ml possess mild to moderate activity of antibacterial and fungal organism when compared to standard drug. (table 1&2)

Table 1: Anti bacterial activity of ethanol extract leaves of *Pseudarthritis viscida* (Linn)

S.NO	ORGANISM	CONCENTRATION OF ETHANOLIC EXTRACT & ZONE OF INHIBITION (mm)				
		25µg/ml	50µg/ml	75µg/ml	100µg/ml	Standard (µg/ml) (gentamycin)
1.	<i>Bacillus megaterium</i>	10	10	12	13	12
2.	<i>E.Coli</i>	--	----	10	12	14
3.	<i>B.Subtilis</i>	11	12	14	16	14
4.	<i>Pseudomonas aeruginosa</i>	--	---	08	10	12
5.	<i>Proteus vulgaris</i>	10	12	12	14	12

Table 2: Anti fungal activity of ethanol extract leaves of *Pseudarthritis viscida* (Linn)

S.NO	ORGANISM	CONCENTRATION OF ETHANOLIC EXTRACT & ZONE OF INHIBITION (mm)				
		25µg/ml	50µg/ml	75µg/ml	100µg/ml	Standard (µg/ml) (fluconazole)
1.	<i>Candida albicans</i>	10	10	12	14	12
2.	<i>Aspergillus Niger</i>	08	11	12	13	12

### CONCLUSION

Medicinal plants are the richest source of drugs of traditional system of medicines and modern medicines, food supplements, folk medicines, nutraceuticals, pharmaceutical intermediates, chemical entities for synthetic drugs. Recent years, ethno botanical, traditional uses of natural compounds specially from plant origin received very much attention as which they are well tested for efficacy, also generally believed to be safe for human use, moreover which is the best classical approach in the search of new molecules for the management of various diseases<sup>11</sup>. The results obtained from the present research indicate ethanol extract of *Pseudarthritis viscida* leaf has potential to act as anti-bacterial and fungal activity and also due to the presence of carbohydrates, flavonoids like active compounds.

### REFERENCES

1. M.Rajina, P.Ratheesh-Chandra, K.M.Khalee, Phytochemical and GC-MS Analysis of leaf extracts of *Pseudarthritis viscida* (Linn) weight & arn. International journal of pharmaceutical sciences and research.2017:8(9):3843-3846.
2. www.flowersofindia&http://indiabiodeversity.org/species.
3. Warrier ps, Indian medicinal plants, orient Longman private limited, new delhi. 1<sup>st</sup> edition 1994.
4. Deepa MA, Narmatha BV and Basker. Antifungal properties of *Pseudarthritis viscida*, fitoterapia, 2004:75(6):581-584.
5. Vijayabaskaran M. Venkiteswaramurthy N. in vitro anti oxidant evaluation of *Pseudarthritis viscida*, international journal of pharmaceutical research.2010:2(3):21-23.
6. Thinagaran rajan, suriyavathana muthukrishnan, characterization of phenolic compounds in *Pseudarthritis viscida* root extract by HPLC and FT-IR analysis, Asian journal of pharmaceutical and clinical research, 2013:6(2):274-276.
7. Hemlal H, Subban Ravi GC-MS, HPTLC antimicrobial analysis of root extracts of *Pseudarthritis viscida* weight and arn and desmodium gangeticum(linn), international research journal of biological sciences.1012:1(5):57-65.
8. Kokate CK, Pharmacognosy, 4th edition, nirali prakashan, New Delhi: 1998: 108-109:127-128.
9. Kavitha K, Nehla Yahoob, Vijayakumar, Reshma Fathima K, Synthesis and Evaluation of Quinazolinone derivatives Asian journal of research chemistry,2017:10(4):577-581.
10. N.Bhaskar, kp. Jaiganesh, R.nepolean, N.selvakumar, evaluation of antimicrobial activity of methanolic extract of *pseudarthritis viscida* linn, international journal of pharmaceutical and chemical sciences.,2012:1(4): 1445-1446.

11. Kavitha K, pharmacological and Phytochemical evidence of Scutia genus plants - a review, Indo American journal of pharmaceutical sciences.2018;5(4): 2957-2963.

**Cite this article as:**

Sangeetha N *et al.* Evaluation of anti microbial activity of ethanol extract of *Pseudarthria viscida* leaf. Int. Res. J. Pharm. 2018;9(6):189-191 <http://dx.doi.org/10.7897/2230-8407.096114>

Source of support: Nil, Conflict of interest: None Declared

Disclaimer: IRJP is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IRJP cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of IRJP editor or editorial board members.