



Review Article

ANTIDIABETIC MEDICINAL PLANTS: A REVIEW

Nabarun Mukhopadhyay¹, Sampath.V², Sameer Pai², U.V.Babu², Richard Lobo^{1*}

¹Department of Pharmacognosy, Manipal College of Pharmaceutical Sciences, Manipal Academy of Higher Education, Manipal- 576104, Karnataka, India

²R&D center, The Himalaya Drug Company, Makali, Bangalore- 562162, Karnataka, India

*Corresponding Author Email: richardlobo73@gmail.com

Article Received on: 25/11/18 Approved for publication: 03/01/19

DOI: 10.7897/2230-8407.100237

ABSTRACT

Diabetes is a metabolic disorder which is mainly characterized by hyperglycaemia and arises by the defects in insulin secretion or insulin action or both. It is categorized into two types, type-I and type-II diabetes. The risk factors that are responsible for diabetes are genetic factor, obesity, hypertension etc. Conventionally many drugs are used for the treatment of diabetes such as biguanides, sulfonylureas, meglitinides etc. But the desired effective treatment is still not to be achieved. So researches are going on for the development of alternate effective therapy against diabetes. Medicinal plants are promising source and also very useful for the development of complimentary therapy. In India medicinal plants are widely used traditionally for the prevention and cure of diabetes. This review article consists the description of the herbs which are reported to have good antidiabetic property.

KEYWORDS: Diabetes, alternative therapy, herbs, hyperglycaemia.

INTRODUCTION

Diabetes mellitus is a very common metabolic disorder which affects the human population throughout the world, characterized by hyperglycaemia and arises due to defects in insulin secretion, insulin action or both. Chronic hyperglycaemia which is a common effect of uncontrolled diabetes causes long-term damage, dysfunction and failure of several organs such as kidneys, eyes, nerves, heart and blood vessels¹. Diabetes is mainly categorized into two types, type-I diabetes and type-II diabetes.

Type-I diabetes is known as insulin dependent diabetes and characterized by deficient production of insulin, requires daily administration of insulin. This happens due to the cellular-mediated autoimmune destruction of the β cells of pancreas.

Type-II diabetes commonly known as non-insulin dependent diabetes which causes due to the ineffective use of the insulin by the body. The following risk factors are commonly involved in the development of type-II diabetes such as genetic factors, obesity, poor diet, insufficient physical activity, advancing age, hypertension etc.²⁻³. There is a another class of diabetes is reported known as gestational diabetes which mainly arises from glucose intolerance, with an onset during pregnancy. This is a temporary condition but it may carries the long term risk of diabetes⁴.

According to World Health Organization, in 2014 it is found that 422 million adults are affected by diabetes. It causes 1.5 million deaths in 2012 and it is seen that the majority of population are mainly affected by type-II diabetes. It is also found that previously type-II diabetes was diagnosed among adults but now it occurs in children also⁵. The oxidative stress is a known

pathogenic mechanism in the development and progression of diabetes which causes due to increased free radical production and impaired antioxidant defenses⁶⁻⁸.

Many drugs are used conventionally for the prevention and management of diabetes such as biguanides, sulfonylureas, meglitinides, PPAR- γ agonists (glitazones), α -glucosidase inhibitors, DPP-4 inhibitors, SGLT2 inhibitors, dopamine-2 agonists etc. But still effective treatment against diabetes yet to be achieved⁹. Research is going on for establishing alternative effective therapies against diabetes. The medicinal plants played an important role in this research as they always are an exemplary source of drugs. In India many herbs are found to be useful for the management of diabetes. From the ethnobotanical information it is found that approximately 800 plants may possess antidiabetic potential¹⁰.

From the knowledge and detailed survey of the medicinal plants we might be able to discover new drugs which are therapeutically active and also cheaper. There are certain advantages of using herbs in the treatment, such as they are easily available, low side effects etc. In this review description about the antidiabetic medicinal plants are given which are reported to have good therapeutic activity by thorough literature survey.

REVIEW

The review work was carried out by thorough searching of different research articles and patents from different online journals from Pubmed, Google scholar, Science direct etc. The literature of scientifically validated plants are also collected which are having good antidiabetic property.

***Pterocarpus santalinus* (Fabaceae)**

The plant is commonly known as red sandalwood and saunderswood is widely found in South India. It contains several chemical constituents, such as carbohydrates, steroids, anthocyanins, saponins, tannins, phenols, triterpenoids, flavonoids, glycosides¹¹. The main active constituent of the plant is santalin. It also contains pterocarpol, pterocartriol, ispterocarpolone, pterocarpo-diolones with β -eudeslone and cryptomeridol¹². It is used as antidiabetic, antihelmintic, aphrodisiac and astringent tonic. It is also used in the treatment of inflammation and ulcer. Several studies are reported which shows about the antidiabetic property of the plant. The heartwood cups for drinking water is the traditional use of the plant in the treatment of diabetes¹³⁻¹⁴. Rao *et al* reported that the hypoglycemic activity at a dose level of 0.25 g/kg body weight/day was showed by the ethanolic fraction of the plant¹⁵. Halim and Mishra found that when the 250 mg/kg of aqueous extract of the plant is coadministered with vitamin E to the streptozotocin induced diabetic rats then it caused significant lowering of blood pressure¹⁶.

***Brassica juncea* (Brassicaceae)**

This plant is commonly known as brown mustard, Chinese mustard is widely found in Europe, Africa, North America and Asia. It contains several important chemical constituents such as triterpenes, Saponins, alkaloids, flavonoids etc¹⁷. It is pharmacologically used as antiscorbutic, diuretic, stimulant, stomachic, antihelmintic, antidysentric, diaphoretic, antiarthritic. Thirumalai T *et al* reported that the aqueous seed extract of the plant showed potent hypoglycemic activity at a dose level of 250, 350 and 450 mg/kg in streptozotocin induced diabetic male albino rat¹⁸.

***Swertia punicea* (Gentianaceae)**

The plant is commonly available in India, Pakistan, China, Japan and other Asian countries. It contains important group of constituents such as xanthonoids, terpenoids, flavonoids, alkaloids, irridoid glycosides etc. Among them xanthonoids are the most abundant class of compounds¹⁹. It is therapeutically used as a hypoglycemic, antihepatotoxic, anti-inflammatory, anti-malarial, antioxidant, antimicrobial etc. There are numerous studies are reported about the antidiabetic property of the herb. A study which is reported by Pen and Fang clearly states that this plant has a potent hypoglycemic effect²⁰. Wen L., Chen J.C reported that the hypoglycemic effect was showed by the ethanol extract and ethyl acetate soluble fractions of the plant in streptozotocin induced type-II diabetic mice²¹. A study reported by L.Y. Tian *et al* states that the significant antidiabetic effect was showed by the xanthone derivatives Methylswertianin and Bellidifolin isolated from the plant at the dose of 200mg/kg body wt. /day in the Streptozotocin (STZ)-induced type 2 diabetic male BABL/c mice²².

***Gymnema sylvestre* (Asclepiadaceae)**

It is a perennial woody vine that is mainly grown in tropical areas of India, Africa and Australia. The main active constituent of the plant is gymnemic acid. Other active constituents are flavones, anthraquinones, henti-acontane, pentatria contane, α and β chlorophylls, phytin, resins, d-quercitol, tartaric acid, formic acid etc²³. Daisy *et al* found that in streptozotocin induced diabetic rats the crude extracts of the plant and its isolated compound dihydroxy gymnemic triacetate showed significant hypoglycemic effect in dose and time dependent manner²⁴. A study reported by Liu *et al* clearly states that insulin secretion from mouse cells was stimulated by the aqueous extract of the leaves of the plant and isolated human islets *in vitro* without

compromising cell viability at the doses of 0.06-2 mg/ml and 0.125 mg/ml²⁵.

***Ricinus communis* (Euphorbiaceae)**

This plant is a species of perennial flowering plant and widely available in India. It contains several chemical constituents such as steroids, Saponins, alkaloids, flavonoids and glycosides etc. The seeds and fruits contain 45% of fixed oil which contains glycosides of ricinoleic, isoricinoleic, stearic, dihydroxystearic acids²⁶. Pharmacologically it is used as an anticancer, antioxidant, antiulcer, antimicrobial, antidiabetic, hepatoprotective, central analgesic, antiasthmatic etc. Dhar ML *et al* reported that the root, stem and leaves of the plant showed hypoglycemic activity in the albino rats at the dose of 250 mg/kg in the initial screening studies²⁷. Poonam Shokeen *et al* reported that the ethanolic extract of the roots of the plant showed potent blood glucose lowering activity in alloxan induced diabetic rats at a dose of 500 mg/kg²⁸.

***Combretum micranthum* (Combretaceae)**

This plant is commonly found in Northwestern Nigeria, Senegal, Mali and Gambia. It contains several classes of compounds such as alkaloids, flavonoids, tannins, terpenoids etc²⁹. It has several pharmacological uses such as antioxidant, anti-inflammatory, antimicrobial etc. Aminu Chika *et al* reported about the potential antidiabetic property of the aqueous leaf extract of the plant at a dose level of 100mg/kg in diabetic and sub-diabetic rats³⁰.

***Arbutus unedo* (Ericaceae)**

It is an evergreen shrub which is native to the France and Ireland. There are different classes of compounds are present in this plant such as terpenoids, α -tocopherol, essential oils, phenolic compounds and organic acids³¹. It is used in the treatment of cardiovascular diseases, gastrointestinal disorders, dermatological problems, diabetes, inflammation, hypertension etc. Mrabti H *et al* reported that the aqueous extract of the plant showed significant *in-vitro* and *in-vivo* antidiabetic effect in the streptozotocin-nicotinamide (STZ-NA) induced diabetic mice at the dose of 500 mg/kg and also it improves metabolic alternations³².

***Cocos nucifera* (Arecaceae)**

It is commonly known as coconut widely available in the coastal regions of India. It contains several important chemical constituents such as phenols, tannins, leucoanthocyanidins, flavonoids, triterpenes, steroids, and alkaloids³³. It is used pharmacologically as an analgesic, anti-inflammatory, antibacterial, antifungal, antioxidant, antiparasitic, antimalarial, cardioprotective etc. S. Saranya *et al* found that the significant antidiabetic activity was showed by the flowers extract of the plant in the streptozotocin induced diabetic rats at the dose of 300 mg/kg³⁴. Nidhi Tyagi *et al* reported that ethanolic extract of this plant showed potent antidiabetic activity at the doses of 200 mg/kg and 400 mg/kg in the streptozotocin induced diabetic rats³⁵.

***Sarcopoterium spinosum* (Rosaceae)**

This plant is commonly found in the Mediterranean region and Middle East. Several flavonoids and phenolics compounds are present in this plant such as Quercetin, hesperidin, naringin, rutin, gallic acid, catechol, protocatechuic acid etc³⁶. It was used traditionally used for the treatment of diabetes. It is also used in the treatment of digestive problems, inflammation and pain. A study reported by Dafni *et al* states that the root extract of the plant was used in muslim folk medicine for the treatment of

diabetes³⁷. Polina Smirin *et al* reported that the root extract of the plant showed potent hypoglycemic activity at a concentration of 0.01 mg/ml in diabetic KK-A^y mice³⁸.

***Vernonia anthelmintica* (Asteraceae)**

The commonly known as wild cumin is an annual herbaceous plant and widely found throughout the India. The main active constituent of the plant is Delta-7-avenasterol that is a steroidal compound which is present in the seeds of the plant. It also contains other constituents such as p-hydroxybenzoyl- vernovan and vernovan which are flavan glycosides, other sterols such as 4-alpha-methylvernosterol, vernosterol, bitter principle demanolide lactone etc³⁹. It is used in the treatment of inflammation, arthritis, diabetes, cancer etc. Shaik Sameena Fatima *et al* reported that the ethanolic extract of the plant showed potent antihyperglycemic activity at a dose of 100 mg/kg in streptozotocin induced diabetic rats⁴⁰.

***Elephantopus scaber* (Asteraceae)**

It is a tropical species of flowering plants which is native to tropical Africa, Eastern Asia, Indian Subcontinent, Southeast Asia, and northern Australia. The major chemical constituents of the plant are sesquiterpene lactones, phenolic acids and flavonoids⁴¹. Pharmacologically it is used as an anticancer, antitumour, anti-inflammatory, antibacterial, antifungal, anticoagulant, hepatoprotective etc. A study reported by Daisy P *et al* clearly states that in streptozotocin induced diabetic rats significant hypoglycemic effect is showed by the acetone extract of the plant⁴².

***Liriope spicata* (Asparagaceae)**

It is an herbaceous flowering plant which is native to East Asia. It contains numerous phytochemical constituents such as steroidal Saponins, polysaccharides etc⁴³. Pharmacologically it is used as an anti-inflammatory, anti-asthmatic, anticancer etc. Xianghong Chen *et al* reported that the aqueous extract of the plant showed significant hypoglycemic effect in streptozotocin (STZ)-induced type 2 diabetic male BABL/c mice at the doses of 200 and 100 mg/kg⁴⁴.

***Mangifera indica* (Anacardiaceae)**

This plant is commonly known as mango which is native to Indian subcontinent. It contains numerous chemical constituents among them polyphenolics, flavonoids and triterpenoids are specific. The main bioactive constituent is mangiferin, which is a xanthone glycoside. The other active constituents present in this plant are isomangiferin, tannins etc⁴⁵. It has several pharmacological uses such as antioxidant, antidiabetic, antiviral, anthelmintic, antiallergenic, antiparasitic, antidiarrhoeal etc. Aderibigbe AO *et al* reported that the aqueous extract of the leaves of the plant showed significant hypoglycemic activity in normoglycaemic and glucose-induced hyperglycaemic mice⁴⁶.

***Aloe vera* (Asphodelaceae)**

It is an evergreen perennial plant which mainly grows in tropical climates around the world and also cultivated in large scale for its medicinal values. This plant mainly contains anthraquinone glycosides such as aloe-emodin, barbaloin, isobarbaloin etc⁴⁷. Ayse Can *et al* reported that aloe leaf gel extract shows beneficial effect as an antidiabetic agent⁴⁸. Jain N *et al* reported that the significant antidiabetic activity was showed by the aloe vera gel at a dose of 200 mg/kg in streptozotocin induced diabetic rats.⁴⁹

***Allium sativum* (Amaryllidaceae)**

This plant commonly known as garlic is native to Central Asia and northeastern Iran and widely found in India. It mainly contains essential oil which consists of sulphur compounds such as trisulfides and disulfides⁵⁰. It is pharmacologically beneficial as an antibacterial, antifungal, antiparasitic, antiviral, antihypertensive, antithrombotic etc. A study reported by Eidi A *et al* states that for the future studies on diabetes mellitus the plant should be considered. Because at the doses of 0.1, 0.25 and 0.5 g/kg in streptozotocin-induced diabetic rats⁵¹.

***Nigella sativa* (Ranunculaceae)**

This plant is also known as black cumin and native to the south and southwest Asia. It contains several chemical constituents such as fixed oil, proteins, alkaloid, saponin, essential oil. The fixed oil contains unsaturated fatty acids such as arachidonic acid, linoleic acid, linolenic acid etc., volatile oil contains saturated fatty acids such as nigellone. The alkaloids are present in this plant are nigellicimine, nigellidine etc⁵². The main active constituent is thymoquinone⁵². It is pharmacologically used as antibacterial, antifungal, antioxidant, antidiabetic, anticancer, immunomodulatory, hepatoprotective etc. Pari L *et al* reported that thymoquinone, the main active constituent of the plant showed significant antidiabetic activity at a dose of 80 mg/kg in streptozotocin-nicotinamide induced diabetic rats⁵³. Najmi A *et al* reported that the *N.sativa* oil showed significant antidiabetic effect in a group of 30 patients. This clinical study was conducted at a tertiary health care center in North India⁵⁴.

***Psidium guajava* (Myrtaceae)**

The plant is commonly known as guava is found widely throughout the India. The leaves of the plant contain an essential oil which is rich in cineol, tannins, triterpenes, flavonoids, resin, eugenol, malic acid etc. The bark contains tannins (12-30%) and calcium oxalate crystals⁵⁵. It is pharmacologically used as an antioxidant, hepatoprotective, anti-allergic, antimicrobial, antiplasmodial, cytotoxic, antispasmodic, cardioactive etc. A study reported by Oh WK *et al* clearly states that the leaf extract of the plant possesses antidiabetic activity in type-II diabetic mice model at a dose of 10 mg/kg⁵⁶.

***Fraxinus excelsior* (Oleaceae)**

This plant commonly known as European ash is widely available throughout the Europe. The chemical constituents present in this plant are iridoids, secoiridoids, flavonoids, triterpenes, alkanes etc⁵⁷. It is used for the treatment of diarrhea, osteoarthritis. Eddouks M *et al* reported that the aqueous extract perfusion of the plant showed significant inhibition in glucose reabsorption in normal and streptozotocin-induced diabetic rats at the dose of 10 mg/kg⁵⁸.

***Cyamopsis tetragonoloba* (Fabaceae)**

It is commonly known as guar or cluster bean is widely cultivated in India. It contains several chemical constituents such as carbohydrates, proteins, fibers, ascorbic acid, flavonoids such as quercetin, kaemferol etc⁵⁹. It is pharmacologically used for the treatment of diabetes, ulcer, hemolysis, asthma, inflammation etc. Mahomed IM *et al* found that the aqueous extract of the plant reduced the blood glucose level in alloxan induced diabetic rats at a dose of 800 mg/kg⁶⁰.

***Origanum vulgare* (Lamiaceae)**

It is a flowering plant which is native to Western and Southwestern Eurasia and also the Mediterranean region. It contains phenolics, phenolic acids, esters and its glycosides, flavonoids, steroids etc⁶¹. It is used pharmacologically as an antioxidant, antifungal, antibacterial, antithrombin etc. Lemhadri A reported that the aqueous extract of the plant showed significant anti-hyperglycemic activity in normal rats at dose of 20 mg/kg⁶².

***Agrimonia eupatoria* (Rosaceae)**

This plant is commonly known as agrimony is available throughout the Europe. It contains many important pharmaceutically active constituents such as terpenes, phenolic compounds, tannins, flavonoids, proteins, carbohydrates and vitamins⁶³. It is therapeutically used as an antiviral, antibacterial, antitumour, analgesic, immunomodulatory, antioxidant, hepatoprotective etc. Gray AM *et al* reported that the aqueous extract of the plant showed significant antihyperglycaemic effect in streptozotocin induced diabetic mice at the doses of 1 mg/ml and 0.25-1 mg/ml⁶⁴.

***Emblica officinalis* (Euphorbiaceae)**

Commonly known as amla, this plant is widely available in India. It contains vitamin C, tannins, gallic acid, ellagic acid, chebullic acid etc⁶⁵. It is used as an antiaging, anti-amnesiac, anti-inflammatory, anti-bacterial, anti-cancer, anti-fungal, anti-viral, anti-venom, anti-ulcerogenic, astringent etc. Shikha Mehta *et al* reported that the aqueous extract of the seeds of the plant showed definite hypoglycaemic activity in streptozotocin induced type-II diabetic rats at the dose of 300 mg/kg⁶⁶.

***Acacia arabica* (Fabaceae)**

It is cultivated throughout the India. It contains phenolics, tannins, gallic acid, protocatechuic acid pyrocatechol, catechin, epigallocatechin-7-gallate, quercetin, isoquercetin etc⁶⁷. It is pharmacologically used as an antibacterial, antifungal, antiviral, antimalarial, antidiarrhoeal, immunomodulatory, anti-inflammatory, antioxidant, antiabortifacient, antiulcer etc. A study reported by Hegazy GA *et al* clearly indicates that this plant extract showed significant hypoglycaemic activity in albino rats at the doses of 100 mg/kg and 200 mg/kg⁶⁸. Mohammad Yasir *et al* found that cold aqueous extract of the plant gives higher antihyperglycaemic activity in diabetic and normal rats at the dose of 400 mg/kg⁶⁹.

***Murraya koenigii* (Rutaceae)**

This plant is commonly known as curry patta is widely available in India. The major chemical constituents present in this plant are α -pinene (51.7%), sabinene (10.5%), β -pinene (9.8%), β -caryophyllene (5.5%), limonene (5.4%), bornyl acetate (1.8%), terpinen-4-ol (1.3%), γ -terpinene (1.2%) and α -humulene (1.2%)⁷⁰. It has several therapeutic activities such as antibacterial, antifungal, antiprotozoal etc. Arulsevan P *et al* found that the plant extract is more effective than glibenclamide in the treatment of diabetes because ethanolic extract of this plant possesses potent hypoglycaemic activity in streptozotocin induced diabetic rats⁷¹. A study reported by S.Vijayanand states that leaf extracts of the plant showed significant hypoglycaemic effect in alloxan induced diabetic albino rats at the doses of 250 and 500 mg/kg⁷².

***Helicteres isora* (Malvaceae)**

This plant is widely distributed throughout the India. It contains important pharmaceutically active chemical constituents such as antioxidants, proteins, phenols, flavonoids, alkaloids, glycosides,

carotenoids, tannins etc⁷³⁻⁷⁴. It is used therapeutically as an expectorant, antidiarrhoeal, antidiabetic, antispasmodic, haemostatic etc. Chakrabarti R *et al* reported that the ethanolic extract of the plant has the potentiality for the treatment of type-II diabetes as it showed significant antidiabetic effect in diabetic C57BL/KsJdb/db mice at the dose of 300 mg/kg⁷⁵.

DISCUSSION

From the ancient times medicinal plants are used for the treatment of many diseases like as diabetes. The herbs are widely used in the treatment because they are considered as more safe and more effective. By utilizing the ethno botanical and ethno pharmacological knowledge we came to know about the medicinal plants which have potent antidiabetic activity. This review article described about the medicinal plants which are commonly used for the treatment of diabetes. The parts of the plant such as leaf and the specific extract such as acetone extract which are more effective are also described here. It is also seen that several researches are also carried out for the desire of establishing an effective treatment against diabetes in India as well as abroad also.

CONCLUSION

In India several medicinal plants are used traditionally in many forms in the treatment of diabetes. Current research of the antidiabetic activity of the herbs helps to develop effective herbal therapies for such purpose. For the discovery of new potential antidiabetic compounds suitable information about medicinal plants are needed. This article is prepared for providing proper information regarding the medicinal plants having antidiabetic property. The informations which are discussed here regarding the medicinal plants might be helpful for further research on diabetes.

ACKNOWLEDGEMENT

The authors like to acknowledge Manipal College of Pharmaceutical Sciences, Manipal Academy of Higher Education, Manipal and The Himalaya Drug Company, Bangalore for their kind support by providing necessary information related to this work.

REFERENCES

1. WHO expert consultation. Report of the expert committee on the diagnosis and classification of diabetes mellitus. *Diabetes Care* 2002; 25: 5–20.
2. Newman B., Selby J. V., King M. C., Slemenda C., Fabsitz, R., Friedman, G. D. Concordance for type 2 (non-insulin-dependent) diabetes mellitus in male twins. *Diabetology* 1987; 30: 763–768. Doi: 10.1007/BF00275741.
3. Kaprio J., Tuomilehto J., Koskenvuo M., Romanov K., Reunanen A., Eriksson J. *et al*. Concordance for type 1 (insulin-dependent) and type 2 (non-insulin dependent) diabetes mellitus in a population-based cohort of twins in Finland. *Diabetology* 1992; 35: 1060–1067. Doi: 10.1007/BF02221682.
4. Bellamy L., Casas J. P. Hingorani, A. D. Williams D. Type 2 diabetes mellitus after gestational diabetes: a systematic review and meta-analysis. *Lancet* 2009; 373: 1773–1779. Doi: 10.1016/S0140-6736(09)60731-5.
5. WHO. Global report on diabetes 2016; Geneva: WHO.
6. Ceriello A. Oxidative stress and glycemic regulation. *Metabolism* 2000; 49: 27–29. Doi: 10.1016/S0026-0495(00)80082-7.
7. Maritim A. C., Sanders R. A., Watkins, J. B. Diabetes, oxidative stress, and antioxidants: a review. *Journal of*

- Biochemical and Molecular Toxicology 2003; 17: 24–38. Doi: 10.1002/jbt.10058.
8. Vos T., Flaxman A. D., Naghavi M., Lozano R., Michaud C., Ezzati M. et al. Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990-2010: a systematic analysis for the global burden of disease study 2010. *Lancet* 2012; 380: 2163–2196. Doi: 10.1016/S0140-6736(12)61729-2.
 9. Inzucchi SE, Bergenstal RM, Buse JB *et al.* Management of hyperglycemia in type 2 diabetes, 2015: A patient-centered approach: Update to a position statement of the American Diabetes Association and the European Association for the study of diabetes. *Diabetes Care* 2015; 38(1): 140-149.
 10. Alarcon-Aguilara FJ, Roman-Ramos R, Perez-Gutierrez S, Aguilar-Contreras A, Contreras-Weber CC, Flores-Saenz JL. Study of the anti-hyperglycemic effect of plants used as antidiabetics. *Journal of Ethnopharmacology* 1998; 61: 101-110.
 11. Narayan S, Devi RS and Devi CSS. Role of *Pterocarpus santalinus* against mitochondrial dysfunction and membrane lipid changes induced by ulcerogens in rat gastric mucosa. *Chemo Biological Interactions* 2007; 170: 67-75.
 12. Yoganarasimhan SN. *Medicinal Plants of India* 2nd edition. New Delhi (India): Cyber media; 2000. p.449.
 13. Nagaraju N, Rao KN. Folk-medicine for diabetes from Rayalaseema of Andhra Pradesh. *Ancient Science of Life* 1989; 9: 31–35.
 14. Nagaraju N, Prasad M, Gopalakrishna G, Rao KN. Blood sugar lowering effect of *Pterocarpus santalinus* (Red Sanders) wood extract in different rat models. *International Journal of Pharmacognosy* 1991; 29: 141–144.
 15. Kameswara Rao B, Giri R, Kesavulu MM, Apparao C. Effect of oral administration of bark extracts of *Pterocarpus santalinus* on blood glucose level in experimental animals. *Journal of Ethnopharmacology* 2001; 74: 69–74.
 16. Halim ME, Misra A. The effect of aqueous extract of *Pterocarpus santalinus* heartwood and vitamin E supplementation in streptozotocin-induced diabetic rats. *Journal of Medicinal Plants Research*. 2011; 5: 398–409.
 17. Parikh H, Khanna A. Pharmacognosy and phytoanalysis of *B. juncea* seeds. *Pharmacognosy Journal* 2014; 6(5): 47-54.
 18. Thirumalai T, Therasa VS, Elumalai EK, David E. Hypoglycemic effect of *Brassica juncea* (seeds) on streptozotocin induced diabetic male albino rat. *Asian Pacific Journal of Tropical Biomedicine* 2011; 4: 323-325.
 19. Goutam Brahmachari, Sadhan Mondal, Arindam Gangopadhyay, Dilip Gorai, Bodhiswatta Mukhopadhyay, Shamal Saha *et al.* *Swertia* (Gentianaceae): Chemical and Pharmacological Aspects. *Chemistry and Biodiversity* 2004; 1: 1627-1651.
 20. Pen F, Fang C.S. Effect of *Swertia punicea* Hemsl. on protection of experimental diabetic mice. *Chinese Journal of Traditional Medical Science and Technol* 2003; 10: 96–97.
 21. Wen L., Chen J.C. Hypoglycemic activity of *Swertia punicea* Hemsl extracts in streptozotocin induced hyperglycemic mice. *China Pharmacy* 2007; 10: 140–142.
 22. L.Y. Tian, X. Bai, X.-H. Chen, J.-B. Fang, S.-H. Liu, J.-C. Chen. Anti-diabetic effect of methylswertianin and bellidifolin from *Swertia punicea* Hemsl. and its potential mechanism. *Phytomedicine* 2010; 17: 533-539.
 23. Gulab S. Thakur, Rohit Sharma, Bhagwan S. Sanodiya, Mukeshwar Pandey, GBKS Prasad, Prakash S. Bisen. *Gymnema sylvestre*: An Alternative Therapeutic Agent for Management of Diabetes. *Journal of Applied Pharmaceutical Science* 2012; 2(12): 1-6.
 24. Pitchai Daisy, James Eliza, Khanzan Abdul, Majeed Mohamed Farook. A novel dihydroxy gymnemic triacetate isolated from *Gymnema sylvestre* possessing normoglycemic and hypolipidemic activity on STZ-induced diabetic rats. *Journal of Ethnopharmacology* 2009; 126: 339–344.
 25. Liu B, Asare-Anane H, Al-Romaiyan A, Huang G, Amiel S.A., Jones P.M., Persaud S.J. Characterisation of the insulinotropic activity of an aqueous extract of *Gymnema Sylvestre* in mouse cells and human islets of langerhans. *Cellular Physiology and Biochemistry* 2009; 23: 125–132.
 26. Manoj Kumar. A Review on Phytochemical Constituents and Pharmacological Activities of *Ricinus communis* L. *Plant. International Journal of Pharmacognosy and Phytochemical Research* 2017; 9(4): 466-472.
 27. Dhar ML, Dhar MM, Dhawan BN, Mehrotra BN, Ray C. Screening of Indian plants for biological activity, Part I. *Indian Journal of Experimental Biology* 1968; 6: 232–247.
 28. Poonam Shokeen, Prachi Anand, Y. Krishna Murali, Vibha Tandon. Antidiabetic activity of 50% ethanolic extracts of *Ricinus communis* and its purified fractions. *Food and Chemical Toxicology* 2008; 46: 3458–3466.
 29. Nounagnon S. Martial, N'tcha Christine, Sina Haziz, Noumavo A. Pacôme, Durand Dah-Nouvlessounon, Assogba M. Fidèle. Antimicrobial activities of *Combretum micranthum* extracts on *Staphylococcus aureus* strains isolated from skin infections and some reference strains. *Asian Journal of Plant Sciences and Research* 2016; 6(4): 40-47.
 30. Aminu Chika, Shaibu Oricha Bello. Antihyperglycaemic activity of aqueous leaf extract of *Combretum micranthum* (Combretaceae) in normal and alloxan-induced diabetic rats. *Journal of Ethnopharmacology* 2010; 129: 34–37.
 31. Ivo Oliveira, Paula Baptista, Albino Bento, José Alberto Pereira. *Arbutus unedo* L. and its benefits on human health. *Journal of Food and Nutritional Research* 2011; 50: 73–85.
 32. Mrabti H, Sayah K, Jaradat, N. et al. (). Antidiabetic and protective effects of the aqueous extract of *Arbutus unedo* L. in streptozotocin-nicotinamide-induced diabetic mice. *Journal of Complementary and Integrative Medicine* 2018; 15(3). Doi: 10.1515/jcim-2017-016.
 33. Lima EBC, Sousa CNS, Meneses LN, Ximenes N.C., Santos Júnior M.A., Vasconcelos G.S. *et al.* *Cocos nucifera* (L.) (Arecaceae): A phytochemical and pharmacological review. *Brazilian Journal of Medical and Biological Research* 2015; 48(11): 953-964. Doi: 10.1590/1414-431X20154773.
 34. S. Saranya, S. Pradeepa. S, S. Subramanian. Biochemical Evaluation of Antidiabetic Activity of *Cocos nucifera* Flowers in STZ Induced Diabetic Rats. *International Journal of Pharmaceutical Sciences Review and Research* 2014; 26(1): 67-75.
 35. Nidhi Tyagi, Vikas Hooda, Anjali Hooda and Sachin Malkani. Evaluation of antidiabetic potential of ethanolic And aqueous extract of *cocos nucifera* endocarp. *World journal of pharmacy and pharmaceutical sciences* 2015; 4(7): 1112-1120.
 36. Seham S El Hawary, Abd El Rahman El Shabrawy, Shahira M Ezzat, Fatma A A El-Shibani. Evaluation of the Phenolic and Flavonoid Contents, Antimicrobial and Cytotoxic Activities of Some Plants Growing in Al Jabal Al-Akhdar in Libya. *International Journal of Pharmacognosy and Phytochemical Research* 2016; 8(7): 1083-1087.
 37. Dafni A., Yaniv Z., Palevitch D. Ethnobotanical survey of medicinal-plants in Northern Israel. *Journal of Ethnopharmacology* 1984; 10: 295–310.
 38. Polina Smirin, Dvir Taler, Guila Abitbol, Tamar Brutman-Barazani, Zohar Kerem, Sanford R. Sampson. *Sarcopoterium spinosum* extract as an antidiabetic agent: In vitro and in vivo study. *Journal of Ethnopharmacology* 2010; 129: 10–17.
 39. Mital N Manvar, T R Desai. *Vernonia anthelmintica* Willd: An Overview on Phytopharmacological Properties. *Inventi Rapid: Ethnopharmacology* 2012; 4:1-4.
 40. Shaik Sameena Fatima, Maddirala Dilip Rajasekhar, Kondeti Vinay Kumar, Mekala Thur Sampath Kumar, Kasetti Ramesh Babu, Chippada Appa Rao. Antidiabetic and antihyperlipidemic activity of ethyl acetate: Isopropanol (1:1)

- fraction of *Vernonia anthelmintica* seeds in Streptozotocin induced diabetic rats. *Food and chemical toxicology* 2010; 48: 495-501.
41. Farha Arakkaveettil Kabeer, Remani Prathapan. Phytopharmacological profile of *Elephantopus scaber*. *Pharmacologia*. 2014; 5(8):272-85.
 42. P. Daisy., R. Jasmine., S. Ignacimuthu., E. Murugan. A novel Steroid from *Elephantopus scaber* L. an Ethnomedicinal plant with antidiabetic activity. *Phytomedicine* 2009; 16: 252-257.
 43. Hung-Jen Lu, Thing-Fong Tzeng, Jui-Chin Hsu, Shih-Hao Kuo, Chia-Hsin Chang, Shin-Ya Huang. An Aqueous-Ethanol Extract of *Liriope spicata* var. *prolifera* Ameliorates Diabetic Nephropathy through Suppression of Renal Inflammation. *Evidence-Based Complementary and Alternative Medicine* 2013; 1-10. <https://doi.org/10.1155/2013/201643>.
 44. Xianghong Chen, Xue Bai, Yihui Liu, Luanyuan Tian, Jianqiu Zhou, Qun Zhou. Anti-diabetic effects of water extract and crude polysaccharides from tuberous root of *Liriope spicata* var. *prolifera* in mice. *Journal of Ethnopharmacology* 2009; 122: 205-209.
 45. Shah KA, Patel MB, Patel RJ, Parmar PK. *Mangifera Indica* (Mango). *Pharmacognosy Reviews*. 2010; 4(7): 42-48. doi:10.4103/0973-7847.65325.
 46. Aderibigbe AO, Emudianughe TS, Lawal BAS. Evaluation of the antidiabetic action of *Mangifera indica* in mice. *Phytotherapy Research* 2001; 15(5): 456-8.
 47. Sharif Moghaddasi M, Sandeep Kumar Verma. *International Journal of Biological & Medical Research* 2011; 2(1): 466-471.
 48. Ayse Can, Nuriye Akev, Nurten Ozsoy, Sehnaz Bolkent, Bahriye Pelin Arda, Refiye Yanardag. Effect of Aloe vera Leaf Gel and Pulp Extracts on the Liver in Type-II Diabetic Rat Models. *Biological and Pharmaceutical Bulletin* 2004; 27(5): 694–698.
 49. Jain N, Vijayaraghavan R, Pant SC, Lomash V, Ali M. Aloe vera gel alleviates cardiotoxicity in streptozocin-induced diabetes in rats. *Journal of Pharmacy and Pharmacology* 2010; 62(1): 115-123.
 50. Lalla fatima Douiri, Ahmed Boughdad, Omar Assobhei, Mohieddine Moumni. Chemical composition and biological activity of *Allium sativum* essential oils against *Callosobruchus maculatus*. *Toxicology And Food Technology* 2013; 3(1): 30-36.
 51. Eidi A, Eidi M, Esmaeili E. Antidiabetic effect of garlic (*Allium sativum* L.) in normal and streptozotocin-induced diabetic rats. *Phytomedicine* 2006; 13 (9-10): 624-629.
 52. Fatemeh Forouzanfar, Bibi Sedigheh Fazly Bazzaz, Hossein Hosseinzadeh. Black cumin (*Nigella sativa*) and its constituent (thymoquinone): a review on antimicrobial effects. *Iranian Journal of Basic Medical Sciences*. 2014; 17(12): 929–938.
 53. Pari L, Sankaranarayanan C. Beneficial effects of thymoquinone on hepatic key enzymes in streptozotocin-nicotinamide induced diabetic rats. *Life Sciences* 2009; 85(23-26): 830-834. Doi: 10.1016/j.lfs.2009.10.021.
 54. Najmi A, Haque SF, Naseeruddin M, Khan RA. Effect of *Nigella sativa* oil on various Clinical and biochemical parameters of metabolic syndrome. *International Journal of Diabetes in Developing Countries* 2008; 16: 85–87.
 55. K. Subashini, Asha P.K. Review for the Phytochemical Constituents Present in *Psidium Guajava* L. and *Psidium Guajava*. *International Journal of Trend in Research and Development* 2017; 4(3): 399-400.
 56. Oh WK, Lee CH, Lee MS, Bae EY, Sohn CB, Oh H. Antidiabetic effects of extracts from *Psidium guajava*. *Journal of Ethnopharmacology* 2005; 96(3): 411-5.
 57. Committee on Herbal Medicinal Products (HMPC). EMA/HMPC/239269/2011. 27 March 2012 Assessment report on *Fraxinus excelsior* L. or *Fraxinus angustifolia* Vahl, folium.
 58. Eddouks M, Maghrani M. Phlorizin-like effect of *Fraxinus excelsior* in normal and diabetic rats. *Journal of Ethnopharmacology* 2004; 94(1): 149-54.
 59. Paras Sharma, Gargee Dubey and Sunil Kaushik. Chemical and Medico-biological profile of *Cyamopsis tetragonoloba* (L) Taub: An overview. *Journal of Applied Pharmaceutical Sciences* 2011; 1(2): 32-37.
 60. Mahomed IM, Ojewole JA. Hypoglycemic effect of *Hypoxis hemerocallidea* corm (African potato) aqueous extract in rats. *Methods and Findings in Experimental and Clinical Pharmacology* 2003; 25(8): 617.
 61. Gottumukkala Venkateswara Rao, T. Mukhopadhyay, T. Annamalai, N. Radhakrishnan, M. R. Sahoo. Chemical constituents and biological studies of *Origanum vulgare* Linn. *Pharmacognosy Research* 2011; 3(2): 143-45.
 62. Lemhadri A, Zeggwagh NA, Maghrani M, Jouad H, Eddouks M. Anti-hyperglycaemic activity of the aqueous extract of *Origanum vulgare* growing wild in Tafilalet region. *Journal of Ethnopharmacology* 2004; 92(2-3): 251-256.
 63. Mikołaj Kostryco, Mirosława Chwil. Biologically active compounds in *Agrimoni eupatoria* L. and their therapeutic effects. *World Scientific News*. 2017; 90-97.
 64. Gray AM, Flatt PR. Actions of traditional anti-diabetic plant, *Agrimoni eupatoria* (*agrimony*): effects on hyperglycaemia, cellular glucose metabolism and insulin secretion. *British Journal of Nutrition* 1998; 80(1): 109-114.
 65. Pushpendra kumar jain, Debajyoti das. Pharmacognostic comparison of *bacopa monnieri*, *cyperus rotundus*, and *emblica officinalis*. *Innovare Journal of Ayurvedic Sciences* 2016; 4(4): 16-26.
 66. Shikha Mehta, Rakesh Kumar Singh, Dolly Jaiswal, Prashant Kumar Rai, and Geeta Watal. Anti-diabetic activity of *Emblca officinalis* in animal models. *Pharmaceutical Biology* 2009; 47(11): 1050–1055.
 67. Mariyam Roqaiya, Wajeeda Begum, Rumaiza Jahufer. *Acacia arabica* (Babool) - A Review on Ethnobotanical and Unani Traditional Uses as well as Phytochemical and Pharmacological Properties. *International Journal of Pharmaceutical and Phytopharmacological Research* 2015; 4 (6): 315-321.
 68. Hegazy GA, Alnoury AM, Gad HG. The role of *Acacia Arabica* extract as an antidiabetic, antihyperlipidemic, and antioxidant in streptozotocin-induced diabetic rats. *Saudi Medical Journal* 2013; 34(7): 727-33.
 69. Mohammad Yasir, Prateek Jain, Debajyoti, M.D.Kharya. Hypoglycemic and antihyperglycemic effect of different extracts of *acacia arabica* lamk bark in normal and alloxan induced diabetic rats. *International Journal of Phytomedicine* 2010; 2: 133-138.
 70. V.S Rana, J.P Juyal, Rashmi, M Amparo Blazquez. Chemical constituents of the volatile oil of *Murraya koenigii* leaves. *International Journal of Aromatherapy* 2004; 14(1): 23-25.
 71. Arulselvan P, Senthilkumar GP, Sathish Kumar D, Subramanian S. Anti-diabetic effect of *Murraya koenigii* leaves on streptozotocin induced diabetic rats. *Pharmazie* 2006; 61(10): 874-877.
 72. Dr.S.Vijayanand. Evaluation of Antidiabetic activity of *Murraya koenigii* on Alloxan Induced Diabetic rats. *International Journal of Pharmaceutical Sciences and Research (IJPSR)* 2015; 6(12); 1401-1405.
 73. Gayathri P., Gayathri Devi, S., Sivagami Srinivasan, S. S. Screening and Quantitation of Phytochemicals and Nutritional Components of the Fruit and Bark of *Helicteres isora*. *Hygeia Journal for Drugs and Medicine* 2010; 2(1): 57-62.
 74. Jain Amita, Sinha Prakriti, Desai Neetin S. Estimation of flavonoid, Phenol content and antioxidant potential of Indian screw tree (*Helicteres isora* L.). *International Journal of*

Pharmaceutical Sciences and Research 2014; 5(4): 1320–1330.

75. Chakrabarti R, Vikramadithyan RK, Mullangi R, Sharma VM, Jagadheshan H, Rao YN et al. Antidiabetic and hypolipidemic activity of *Helicteres isora* in animal models. *Journal of Ethnopharmacology* 2002; 81: 343-349.

Cite this article as:

Nabarun Mukhopadhyay *et al.* Antidiabetic medicinal plants: A review. *Int. Res. J. Pharm.* 2019;10(2):31-37
<http://dx.doi.org/10.7897/2230-8407.100237>

Source of support: Nil, Conflict of interest: None Declared

Disclaimer: IRJP is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IRJP cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of IRJP editor or editorial board members.